

RESEARCH

Open Access

Spatial heterogeneity of urban soils: the case of the Beijing metropolitan region, China

Qizheng Mao¹, Ganlin Huang¹, Alexander Buyantuev², Jianguo Wu^{1,3}, Shanghua Luo⁴ and Keming Ma^{5*}

Abstract

Introduction: Urban soils are large pools of carbon, nitrogen, and other elements, supporting plant growth, sustaining biogeochemical cycles, and serving as the foundation for maintaining ecosystem function and services of urban green spaces (UGS). Quantifying urban soil properties is essential for assessing urban ecosystem services and detecting pollution. Characterizing spatial heterogeneity of urban soil properties, which may change with land use or urbanization, is crucial for understanding urban ecosystem functions.

Methods: We collected 466 soil composite samples across the Beijing metropolitan region and then analyzed the heterogeneity of soil properties in UGS of Beijing metropolitan region by ANOVA along the urbanization gradient spanning different land uses.

Results: Our results show that soil properties vary considerably among land-use types and along the urban-rural gradient. Soil moisture content, soil organic carbon (SOC), available phosphorus, available potassium, total nitrogen (TN), the ratio of SOC:TN (C/N), Pb, and Cu increased notably from suburbs to the urban core, while soil pH, bulk density, TN, Cr, Ni, and Mn were not significantly different across urbanization levels. Most soil properties, except soil TN, Cu, Cr, Ni, exhibited significant differences between different land uses. The highest levels of soil nutrients and heavy metals were found in roadsides and residential areas among all the land uses.

Conclusions: The spatial heterogeneity of urban soils in the Beijing metropolitan region is mainly attributable to the different land uses. Soil nutrient content has increased in UGS due to fertilization and pollution. Contamination of urban soils by heavy metals is due largely to traffic emissions and pollution by industrial and household wastes. Although human activities tend to augment urban soil nutrient pools to maximize certain ecosystem services, they may inadvertently compromise these and other services by increasing pollution. Knowledge of urban soil spatial heterogeneity in UGS is indispensable for improving urban ecosystem management.

Keywords: Urbanization; Urban green spaces; Soil properties; Land use; Urban ecology; Beijing

Introduction

Soil, as the component of urban green spaces (UGS) (Setälä et al. 2013), plays a pivotal role in maintaining urban ecosystem services, such as biodiversity maintenance, water resources protection, microclimate regulation, carbon sequestration, food production, and fulfillment of cultural or recreational needs (Lovell and Taylor 2013; Neil et al. 2014; Wu 2014). Urban soils are large pools of carbon, nitrogen, and other elements, serving as the foundation of urban ecosystems by supporting plant growth

and sustaining biogeochemical cycles (Pouyat et al. 2002). Quantifying soil properties and understanding soil conditions are essential for assessing ecosystem services provided by UGS and detecting pollution in UGS.

Urban soils are often physically disturbed, chemically polluted, and tend to be highly heterogeneous spatially (Wu 2014). Previous studies conducted at regional scales examined soil properties' variations along urbanization gradients of Hong Kong (Jim 1998a), Phoenix (Zhu et al. 2006), Baltimore, and New York (Pouyat et al. 2008). Major findings from these and other studies are as follows. First, urbanization is accompanied by the transporting of large amounts of rock material in the landscape, the introduction of construction rubble, bricks, and other building

* Correspondence: mkm@cees.ac.cn

⁵State Key Laboratory of Urban & Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, P. R. China

Full list of author information is available at the end of the article

materials (Jin et al. 2011). Consequently, urban soils often are compacted and structurally degraded, alkaline, and water deficient (Jim 1998a), which makes them less suitable for growing plants. Second, existing soil patterns are the product of both human disturbances and management-induced amelioration and therefore the effects of urbanization on soil nutrients. For example, some researcher demonstrated that urban soil had lower nutrients (Jim 1998b; Baxter et al. 2002), while other studies pointed out that soil nutrients were higher in urban area (Kaye et al. 2008) and had sufficient nutrient levels to support plant growth (Pouyat et al. 2007). Third, pollutants from industrial activities (Shang et al. 2012), heavy traffic (Chen et al. 2010a; Hamzeh et al. 2011), and house wastes (Schwarz et al. 2012; Szolnoki et al. 2013) all contribute to soil contamination by heavy metals, which is likely to have adverse effects on human health.

At local scales, urban soil properties were compared between different land uses in a number of cities, including Baltimore (Pouyat et al. 2007), Hong Kong (Jim 1998a), Hubei (Li et al. 2013), and Lanzhou (Wang et al. 2011). It is widely agreed that the difference in human activities contribute to the variation of soil properties among land-use types (Pouyat et al. 2007; Davies and Hall 2010). For example, soils in transportation and residential areas were characterized by higher pH and bulk density (BD) (Pouyat et al. 2007; Zhao et al. 2013; Yang et al. 2014). Soils in urban forests (Zhao et al. 2013) or protected places (Yang et al. 2014) contained higher soil organic carbon (SOC), while soils at industrial sites and in transportation areas were often polluted by heavy metals (Lu and Bai 2010; Hamzeh et al. 2011). At finer scales, urban soil carbon and nitrogen also varied within single patches, although this variation was smaller than that at broader scales and different land uses (Jenerette et al. 2006).

Previous investigations undoubtedly suggest significant transformations in urban soil properties as a result of land use and land cover changes. However, a deep understanding of the urban soil biogeochemistry is still limited, requiring more research to be conducted. Considerable spatial heterogeneity of urban soils warrants a landscape ecological perspective for analyzing patterns and determinants of soils in urban areas. We therefore investigated urban soil variation in UGS of Beijing along the urbanization gradient and across different land-use types in order to understand this heterogeneity. Specifically, we focused on three questions: (1) What are the soil properties in UGS in Beijing and do they meet the needs of plant growth or not? (2) How and do soil properties vary along urbanization gradients or not? (3) How and do soil properties vary across land-use types or not? Our findings are expected to have implications for a better understanding of urbanization effects on soils and contribute to scientifically informed soil management and conservation in UGS.

Methods

Study area

Beijing, one of the oldest cities in the world which has experienced rapid urbanization in the recent decades, was chosen for our case study. It is located in the northeast of the North China Plain and has an area of approximately 16,410 km² and a population of 20.69 million (Beijing Municipal Bureau of Statistics 2013). During 1984 and 2008, the urbanized area in Beijing increased 3.4 times (Kuang 2012), while permanent residents increased by 53.4% (Beijing Municipal Bureau of Statistics 2013). Beijing has diverse land uses including residential (52.64%), wood (7.04%), institutional (6.83%), industrial, and commercial (4.62%), transportation (2.91%), and others (Kuang 2012). Such degree of urbanization and variability of land-use types makes Beijing a good study site to address our research questions.

Soil sampling

In July 2009, we set up eight transects from the urban core to suburban areas in eight directions (east, west, north, south, northeast, northwest, southeast, southwest) in order to capture urbanization gradients. Along these transects, 78 survey plots spaced at 3-km interval were selected. Another 52 plots at 6-km interval were later added to densify our sampling network and capture more of the spatial heterogeneity. Consequently, 130 plots were sampled (Figure 1). We recorded dominant land use within the 500-m radius of each plot. One composite surface (0 to 20 cm) soil core was obtained within a representative 20 × 20-m open space by mixing three cater-cornered subsamples. Three additional samples were collected by cutting rings and then transferring them into three aluminum boxes. A total of 466 soil composite samples were collected during July to September.

Sample preparation and instrument analysis

Soil samples were stored indoor and dried at room temperature. Debris (e.g., leaves, stones, worms) were handpicked and removed. The samples were then grounded to go through 2-, 1-, and 0.15-mm nylon sieves, respectively. Laboratory analyses for soil properties were then conducted using soil composites through different nylon sieves. Specifically, we analyzed the soil moisture content (SMC), soil pH, soil BD, total nitrogen (TN) content, SOC:TN (C/N), SOC, available phosphorus (AP), available potassium (AK), and heavy metals (Pb, Cr, Cu, Ni, Mn).

All laboratory tests followed standard methods. Using soils from the aluminum boxes, SMC was determined by drying them at 105°C for 24 h and was calculated as the percentage of water in oven-dry soil (%). BD was determined from the total oven-dry mass and was calculated as milligrams of oven-dry soil in the volume of 1 cm³. Using soils through 2-mm nylon sieves, pH was measured with a 1:2.5 mixture




Figure 1 Study area and soil sampling design.

(soil:water ratio) by the glass electrode method (McLean 1982). Using soils through 1-mm nylon sieves, we measured AP by Mo-Sb colorimetry from a sample extracted with NaHCO_3 and non-phosphorus active carbon (Tiessen et al. 1993), while AK was extracted with ammonium acetate and examined directly by inductively coupled plasma optical emission spectroscopy (ICP-OES) (Simard 1993). Using soils through 0.15-mm nylon sieves, TN and TC were analyzed by the dry combustion method (Vario EL Element Analyzer) (Bremner and Mulvaney 1982). SOC was measured by potassium dichromate titration method (Lu 2000). Finally, we measured the contents of Pb, Cu, Cr, Ni, and Mn by extracting them using HNO_3 , HClO_4 , and HF according to USEPA method 3052 with a microwave laboratory unit (USEPA United States Environmental Protection Agency 1995). Pb, Cr, Mn, Cu, and Ni contents were all determined using ICP-OES.

Urbanization gradient and classification of land use

Beijing has five circular express roads, which are named as the second to the sixth ring road. The rapid urban sprawl in Beijing followed a homocentric expansion along this ring-road system (Kuang 2012). We therefore use these ring roads as benchmarks of urbanization phases in Beijing. According to this logic, we define the area within the third ring road as the urban core area, while the area between the third and the fifth ring roads as the urban transitional zone. The suburban area encompasses developments outside the fifth ring road (Figure 1).

We categorized all 466 samples to seven classes according to the dominant land use and land cover type (Figure 2), and they were as follows:

- (A) Park ($n = 57$), open space for recreational use
- (B) Residential area ($n = 78$), residential buildings
- (C) Institutional area ($n = 40$), school, hospital, and office places
- (D) Road greenbelts ($n = 114$), large areas of ribbon vegetation distributed along roadsides
- (E) Roadside ($n = 104$), single tree rows along major roads
- (F) Industrial area ($n = 17$), factories
- (G) Forest ($n = 57$), large area of vegetation and plants in suburban areas

Statistical analyses

Statistical analyses were performed in SPSS 21. The Kolmogorov-Smirnov (K-S) test for normality was used to determine whether variables were normal or skewed. Data were log transformed to stabilize the variance of soil properties for AP, AK, SOC, TN, C/N, Pb, Cu, Ni, and Mn. The analysis of variance (ANOVA) was used to test whether soil properties varied between different areas followed by multiple comparisons to test the effect of land-use types and urbanization on each variable. To deal with sample size inequalities for different land-use types or urbanization levels, we employed the Hochberg's method for *post hoc* analysis to determine significant differences between mean soil variables (Pouyat et al. 2007).




Figure 2 Seven types of land use in our study. (a) Road greenbelts, (b) roadside, (c) residential area, (d) neighborhood park, (e) forest, (f) industrial area, (g) institutional sites.

Background values of soil properties in Beijing

To articulate human activities' impacts on UGS soil properties, we compared our samples with background values of soil properties in Beijing (Su et al. 2001; Liu et al. 2003), which were sampled in mountains in the outskirts of Beijing to represent soil properties values without human disturbance. We refer to China's Environmental Protection Agency's work for these background soil values (China National Environmental Monitoring Center 1990).

Results

General characteristics of urban soils in Beijing's UGS

Descriptive statistics of each soil variable revealed that soil properties in UGS varied widely (Table 1). The highest coefficient of variation (CV) was found in soil AP (119.25%) and the lowest in soil pH (2.36%).

Comparing our results with soil background values, we found that the mean values of soil pH, BD, and AP were higher, AK, SOC, TN, and C/N were lower, and SMC was similar to that in Dongling mountain in

Table 1 Descriptive statistics of soil properties in Beijing

Soil properties	Median	Minimum	Maximum	SE	CV (%)	Mean	Background value	K-S P value
pH	8.07	7.43	8.77	0.01	2.36	8.05	6.47 ^a	0.010
BD (mg/kg)	1.39	0.89	1.74	0.01	9.13	1.38	1.11 ^b	0.054
Moisture (%)	17.35	3.79	54.84	0.28	34.90	17.02	16.62 ^a	0.278
AP (mg/kg)	14.33	0.61	274.03	1.16	119.25	20.71	11.37 ^a	<0.001
AK (mg/kg)	158.13	41.20	711.10	4.16	50.54	176.44	281.8 ^a	<0.001
SOC (g/kg)	9.61	1.68	64.28	0.26	53.94	10.51	62.16 ^a	<0.001
TN (g/kg)	0.85	0.23	2.46	0.01	33.91	0.89	2.84 ^a	0.009
C/N	11.16	4.08	26.13	0.15	27.77	11.59	21.54 ^a	<0.001
Pb (mg/kg)	18.53	0.66	291.98	0.83	81.45	21.80	25.40 ^c	<0.001
Cr (mg/kg)	54.11	5.24	154.75	0.54	21.08	54.14	68.10 ^c	<0.001
Cu (mg/kg)	25.15	0.70	125.22	0.64	47.65	28.39	23.60 ^c	<0.001
Ni (mg/kg)	25.03	11.84	61.56	0.21	18.02	25.33	29.00 ^c	0.010
Mn (mg/kg)	672.10	338.21	1572.52	5.76	17.84	685.49	705.00 ^c	0.049

^aLiu et al. 2003, a study of soil properties in mountain region, Beijing; ^bSu et al. 2001, another study of soil properties in mountain region, Beijing; ^cChina National Environmental Monitoring Center 1990, a book of soil background value in China. SE-Standard errors; CV-Coefficient of variation.

Beijing. Of these, only mean Cu was higher than the soil background value of Beijing (23.6 mg/kg), which signifies relatively clean soils, close to natural soils.

Variation of soil properties along urbanization gradients

Table 2 shows variation of soil properties along the urbanization gradient. Soil moisture, AP, AK, SOC, C/N, Pb, and Cu showed significant increase from suburban area to the urban core (Figure 3). The SMC in suburban areas (16.25%) is not statistically different ($P = 0.250$) from the transitional zone (17.4%) but both are significantly different from the urban core (20.12%). Soil AP and AK were significantly different between different urbanization levels ($P < 0.001$), but the difference of SOC was only detected between the suburban area and the urban core ($P = 0.012$), while the difference of soil C/N was detected between suburban area and other two areas. The soil Pb was significantly higher ($P < 0.001$) in the urban core and the transitional zone compared to suburban areas, whereas no significant difference of soil Pb is detected between the urban core and the transitional zone ($P = 0.659$). Soil Cu content was significantly higher in the urban core area when compared to the transitional zone and suburban areas. For soil pH, BD, TN, Cr, Ni and Mn, no significant difference was found along the urbanization gradient.

Variation of soil properties within different land-use types

The soil properties were notably different between land-use types (Table 3). Our results showed that soil pH, BD, SMC, AP, AK, SOC, C/N, TN, Pb, and Mn were significantly different between the seven land-use types ($P < 0.05$). Detailed results on how each soil property varies

across land-use types are listed in Table 3. In particular, we want to point out the following soil property variations across land-use types and will discuss what caused such variation in the 'Discussion' section.

The soil pH, BD, and AP in roadsides and residential areas are significantly higher than in the forests. The soil Mn, SOC, and C/N are higher in roadsides than the rest land-use types. Soil AP is significantly higher in road greenbelts.

Discussion

Soil conditions of UGS in Beijing

Plant growth


Our results indicate that soils are practically unsuitable for plant growth in many places within the Beijing metropolitan region. The soils of UGS in this region are commonly alkaline and mechanically compacted. The soil pH in all sites exceeds the pH level of neutral soils (pH = 7), which makes them unsuitable for growing most of plants according to earlier recommendations (Whitcomb 1987). The high levels of alkaline can influence soil nutrient solubility (Messenger 1986) and lead to soil fertility imbalance, deficiency, and chlorosis symptoms (Harrell et al. 1988). Furthermore, measured soil BD was above 1.10 mg/m³ for 98.92% of the samples and 1.49 mg/m³ for 18.2% of the samples, which are expected to affect plant root growth (USDA United States Department of Agriculture 2001). This is also aggravated by reduced soil moisture holding capacity, aeration, and drainage.

Our results also suggest deficiencies in soil AP and SOC. While soil AP in urban sites is almost twice as high (11.67 mg/kg) as the background value (Liu et al. 2003), it is still lower (less than 30 mg/kg in 82.17% of

Table 2 Mean (\pm SE) surface soil properties (0 to 20 cm) along the urbanization gradient

Soil properties	Suburban area (n = 305)	Transitional zone (n = 98)	Urban core (n = 63)	Background value	P value
pH	8.03 \pm 0.01	8.07 \pm 0.02	8.09 \pm 0.02	6.47	0.076
BD (mg/m ³)	1.38 \pm 0.01	1.39 \pm 0.01	1.39 \pm 0.02	1.11	0.836
SMC (%)	16.25 \pm 0.33 ^a	17.4 \pm 0.6 ^a	20.12 \pm 0.79 ^b	16.62	<0.001
AP (mg/kg)	16.08 \pm 1.31 ^a	22.09 \pm 2 ^b	40.25 \pm 3.9 ^c	11.37	<0.001
AK (mg/kg)	153.61 \pm 4.21 ^a	188.83 \pm 7.61 ^b	266.65 \pm 14.66 ^c	281.8	<0.001
SOC (g/kg)	10.19 \pm 0.35 ^a	10.76 \pm 0.52 ^{ab}	11.68 \pm 0.54 ^b	62.16	0.012
TN (mg/kg)	0.89 \pm 0.01	0.86 \pm 0.03	0.97 \pm 0.05	2.84	0.09
C/N	11.24 \pm 0.2 ^a	12.16 \pm 0.3 ^b	12.33 \pm 0.34 ^b	21.54	0.001
Pb (mg/kg)	18.84 \pm 1.05 ^a	26.63 \pm 1.57 ^b	29.56 \pm 2.05 ^b	25.4	<0.001
Cr (mg/kg)	53.51 \pm 0.68	56.37 \pm 1.34	53.85 \pm 0.69	68.1	0.074
Cu (mg/kg)	25.78 \pm 0.59 ^a	28.98 \pm 1.59 ^a	39.95 \pm 2.32 ^b	18.7	<0.001
Ni (mg/kg)	25.25 \pm 0.26	25.7 \pm 0.59	25.2 \pm 0.32	29	0.77
Mn (mg/kg)	686.58 \pm 7.87	674.28 \pm 9.98	697.09 \pm 10.86	705	0.43

Values with the same letters in superscript are not significantly different at 5% level based on Hochberg's multiple comparison tests. Last column shows P values for ANOVA tests.



the samples) than the recommended suitability range of 30 to 60 mg/kg for healthy plant growth (Whitcomb 1987). The SOC in the urban area (10.51 g/kg) is more than five times less than the background value (62.16 g/kg) (Liu et al. 2003). Moreover, only 42.25% of the samples are above the recommended range of soil SOC (>10 g/kg) for healthy plant growing (Whitcomb 1987). Such SOC deficiency in UGS can easily deplete soil structural composition and complicate maintenance (Jim 1998b).

In contrast to soil AP and SOC, AK is found in sufficient, even excessive, quantities in Beijing (176.44 mg/kg), which is higher than the recommended range of 78 to 156 mg/kg for healthy plant growth (Whitcomb 1987). The negative side of very high AK in about 50%

of the samples is the potential imbalance in nutrients, which has consequences for Ca absorption (Hamberg et al. 2009; Moussa et al. 2009).

Pollution

The overall health risk of heavy metal pollution in soils of green spaces in Beijing is relatively low whereas some higher localized soil Pb and Cu content exists. Cu in 18.54% of the samples, Pb in 11.23% of the samples, and Cr and Ni in 0.66% of the samples exceed the environmental quality standards in China (35 mg/kg for Cu and Pb, 90 mg/kg for Cr, and 40 mg/kg for Ni). Two out of the 466 samples of Cu and one sample of Ni exceed the second level (100 mg/kg for Cu and 60 mg/kg for Ni). This level corresponds to the safety threshold for human

Table 3 Mean (\pm SE) surface soil properties (0 to 20 cm) of different land uses

Soil properties	Industrial	Roadside	Residential	Institutional	Road greenbelts	Park	Forest	P value
pH	8.02 \pm 0.04 ab	8.12 \pm 0.02 a	8.08 \pm 0.02 a	8.10 \pm 0.02 a	8.05 \pm 0.02 a	7.96 \pm 0.03 b	7.95 \pm 0.02 b	<0.001
BD (mg/m ³)	1.32 \pm 0.04 ab	1.43 \pm 0.01 a	1.42 \pm 0.01 a	1.41 \pm 0.02 a	1.38 \pm 0.01 a	1.38 \pm 0.01 a	1.28 \pm 0.02 b	<0.001
SMC (%)	19.09 \pm 0.84 ac	13.01 \pm 0.66 b	19.50 \pm 0.46 a	18.71 \pm 0.67 ac	18.11 \pm 0.57 ac	17.77 \pm 0.62 ac	16.25 \pm 0.74 c	<0.001
AP (mg/kg)	31.15 \pm 17.54 ab	19.33 \pm 2.30 ab	23.34 \pm 2.04 a	18.48 \pm 2.58 ab	21.79 \pm 1.87 a	20.52 \pm 3.9 ab	16.15 \pm 3.28 b	0.003
AK (mg/kg)	145.36 \pm 16.85 ab	209.76 \pm 9.98 a	206.93 \pm 12.42 a	149.84 \pm 9.93 b	163.49 \pm 6.37 b	137.17 \pm 7.1 b	166.14 \pm 12.48 b	<0.001
SOC (g/kg)	15.04 \pm 4.18 ab	13.09 \pm 0.64 a	10.22 \pm 0.43 b	8.52 \pm 0.49 b	9.46 \pm 0.44 b	9.59 \pm 0.64 b	9.57 \pm 0.49 b	<0.001
TN (g/kg)	1.07 \pm 0.15	0.93 \pm 0.03	0.88 \pm 0.02	0.81 \pm 0.03	0.84 \pm 0.03	0.91 \pm 0.05	0.93 \pm 0.04	0.052
C/N	12.02 \pm 1.06 ab	13.93 \pm 0.37 b	11.39 \pm 0.3 a	10.71 \pm 0.47 a	11 \pm 0.25 a	10.47 \pm 0.33 a	10.24 \pm 0.3 a	<0.001
Pb (mg/kg)	41.51 \pm 18.25 ab	24.74 \pm 1.51 a	23.90 \pm 1.70 ab	20.46 \pm 1.89 ab	19.11 \pm 0.97 b	16.58 \pm 0.9 b	19.79 \pm 1.17 ab	0.010
Cr (mg/kg)	54.03 \pm 2.36	55.56 \pm 1.42	55.11 \pm 1.07	51.99 \pm 2.16	52.64 \pm 1.04	53.69 \pm 1.54	55.19 \pm 0.95	0.279
Cu (mg/kg)	28.82 \pm 4.28	31.63 \pm 1.66	30.17 \pm 1.54	25.71 \pm 1.47	26.76 \pm 1.09	27.39 \pm 2.10	26.12 \pm 1.25	0.068
Ni (mg/kg)	25.40 \pm 1.33	24.35 \pm 0.53	26.13 \pm 0.43	25.30 \pm 0.61	25.15 \pm 0.30	25.61 \pm 0.84	26.11 \pm 0.63	0.071
Mn (mg/kg)	708.08 \pm 73.56 ab	721.64 \pm 11.83 a	683.78 \pm 11.33 ab	677.10 \pm 17.58 ab	673.68 \pm 9.19 ab	681.71 \pm 18.16 ab	646.30 \pm 17.45 b	0.004

Values with same letters are not significantly different at 5% level based on Hochberg's multiple comparison tests. Last column shows P values for ANOVA tests.

health (SEPA China State Environmental Protection Administration of China 1995).

Effects of urbanization on soil properties

Our findings indicate that soil properties change considerably along urbanization gradients, which is consistent with previous researches (Jim 1998a; Zhu et al. 2006; Kaye et al. 2008; Pouyat et al. 2008) (Figure 3). Human disturbance impacts soil properties in UGS mostly through intensive management and waste discharge.

The urban core has significantly higher SMC than the transitional zone and suburban area because UGS in the urban core are intensively maintained and frequently irrigated. Our finding of the increased SMC in the urban core of Beijing echoed the patterns of Phoenix in Arizona, USA (Zhu et al. 2006).

Frequent irrigation and fertilization in urban areas also caused dramatic increase in soil AP, AK, and SOC from suburbs to the urban core (Figure 3), which is consistent with the higher content of soil nutrients in urban soils found previously (Zhu et al. 2006; Kaye et al. 2008; Pouyat et al. 2008, 2010). Direct inputs of pollutants also contribute to such patterns (Kaye et al. 2008). For example, untreated sewage, domestic waste, and excessive fertilizer applications could all lead to AP accumulation in urban and suburban soils (Huang et al. 2013). The urban construction, which uses cement and concrete with abundant Ca, is a source of higher surface charge densities and cation exchange interaction. Its most direct consequence is the fixation of absorbed K ions from soil colloids (Li et al. 2013; Moussa et al. 2009).

Our results show that SOC is positively correlated with the soil Pb ($R^2 = 0.41$, $P < 0.001$) and Cu ($R^2 = 0.18$, $P < 0.001$) (Figure 4). This finding is consistent with what researchers found previously that soil organic matter has a

positive contribution to heavy metal accumulation (Sauvé et al. 2000).

Soil pollution by heavy metals (i.e., Pb and Cu) increased significantly along the urbanization gradient (Table 2). The higher Pb concentrations in the urban core are explained by traffic emission and historical use of leaded paints (Chen et al. 2010b; Xia et al. 2011). Mao et al. (2014) presented an in-depth analysis on spatial heterogeneity of Pb in Beijing UGS.

Three factors contributed to the increase of soil Cu from the suburbs (25.78 mg/kg) to the urban core (39.95 mg/kg) (Figure 3). First, abundant historical sites (temples, palaces, ancient parks) in central Beijing contain many brass cultural relics. Second, Cu is known to be emitted as a result of deterioration of some mechanical parts in vehicles over time. In particular, the lining in automobile brakes was found to be the source of Cu (Hamzeh et al. 2011). Third, application of copper-rich pesticides in UGS may result in an increase of soil Cu in the urban core (Szolnoki et al. 2013).

Effects of land use on soil properties

Most soil properties varied among seven land-use types, and these variations are due mainly to human-induced disturbances, as well as management practices that aim to improve soil conditions. Our results suggest that soils in roadsides and residential areas were most mechanically disturbed and had higher soil nutrients and heavy metal pollutants when compared to other land uses. This finding is consistent with previous findings (Pouyat et al. 2007; Raciti et al. 2008; Davies and Hall 2010; Li et al. 2013).

Roadsides and residential areas

Close to both streets and pedestrians, roadsides and residential areas are exposed to intense human disturbance,




Figure 4 Correlation of soil Pb/Cu with SOC.

such as trampling, sewage discharge, and pet litter (Li et al. 2013; Yang et al. 2014). In contrast, forests in our study are located in infrequently disturbed suburbs away from busy streets and sidewalks (Figure 2e). As a result, soil pH and BD in roadsides and residential areas are significantly higher than in forests.

Roadside and residential areas have higher soil AK ($P < 0.05$). Researchers found that the high level of Ca will increase the level of extractive K and such process is catalyzed by high soil pH (Li et al. 2013). Construction materials (i.e., bricks and cement) with high Ca content are a major component in roadside and residential areas. Therefore, we speculate that the high K in roadside and residential areas is related to the interaction between K and Ca in an alkaline environment. Soil AP is significantly higher in residential areas, which is mainly due to reclaimed water irrigation in these land uses. Roadsides have the lowest SMC. We attribute this to three factors. First, roadsides are less irrigated than UGS in other land uses. Second, proximity to roads and heavy automobile traffic can increase the ambient temperature and soil evaporation (Figure 2b). Third, higher soil BD decreases soil permeability and water infiltration thus increasing runoff.

Roadsides have significantly higher SOC, which can be explained by the following. First, soils in roadsides are seldom disturbed, while other land uses undergo transformations and replacement of vegetation by structures or other vegetation, which depletes SOC. Second, atmospheric deposition from vehicles and high accumulation of dust from nearby sealed surfaces result in increased nutrient content in roadsides (Chon et al. 1998; Park et al. 2010; Li et al. 2013). Third, street litter and animal waste in roadsides may also lead to SOC accumulation (Park et al. 2010; Zhao et al. 2013). Similar to SOC, the highest C/N is found in roadsides followed by industrial areas. However, soil TN shows no statistical difference ($P = 0.052$) between seven land-use types.

Soil Mn is also higher in roadsides. Li et al. (2013) also found highest soil Mn concentrations in this land-use type. The pattern can be explained by higher soil temperature near paved surfaces, which promotes soil mineralization and results in higher nutrient content (Curtin et al. 2011).

Road greenbelts

Soil AP is significantly higher in road greenbelts. We found parts of road greenbelts being irrigated by a mixture of reclaimed water from various urban sewages rich in phosphorous. The highest AP content (274.03 mg/kg) was detected near the steel production factory, the Shougang Construction Group. However, soil Pb and SOC in road greenbelts are relatively low, which is consistent with previous findings (Zhao et al. 2013). This is because most of the road greenbelts in Beijing were newly built and filled with garden soils and have only

experienced a short-time accumulation of soil organic matters and pollutants.

Limitations

Due to limited time and resources, we only measured selected soil properties, while other essential indicators of soil health, such as available nitrogen, micronutrients (e.g., Ca, Mg, Fe), and other heavy metal pollutants (e.g., Cd, Zn, As), were not assessed.

Conclusions

Urban soils in the Beijing metropolitan region are considerably alkaline and compacted. Soil TN, SOC, and AP are in deficit, while AK is abundant and sufficient for supporting plant growing. Heavy metal pollution in Beijing is low. Analyses of the spatial patterns of soil properties in Beijing suggest that soil AP, AK, SOC, C/N, Pb, and Cu increase from suburbs to the urban core, while other elements showed no significant difference. Land use was found as a strong predictor of spatial heterogeneity of soil properties. Roadsides and residential areas are the two land uses characterized by higher soil nutrients and heavy metal pollutants. The major factor of soil spatial heterogeneity in our study area is the time of soil accumulation followed by human activities (e.g., physical disturbance, pollutant emissions, irrigation, fertilization, and soil restoration). Our study is an important step toward understanding the spatiotemporal pattern of soil properties in urban regions. Future research should systematically evaluate both the negative and positive effects of management practices on urban soils and help design effective management plans for ameliorating soil conditions. Also, it is critically important to enumerate the different ecosystem services (and disservices) provided by urban soils.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

QM and KM designed the research; QM and SL conducted the research and analyzed the data; QM drafted the manuscript; GH, AB, and JW revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This research was supported by two projects - the 'Special Foundation of State Key Lab of Urban and Regional Ecology (grant SKLURE2013-1-01)' and the 'Foundation for The Recruitment Program of Global Experts of Beijing Normal University (257-107023)'. The authors sincerely thank Chunyang He, Deyong Yu, and Zexiang Sun for their valuable input.

Author details

¹Center for Human-Environment System Sustainability (CHESS), State Key Laboratory of Earth Surface Processes and Resource Ecology (ESPRE), Beijing Normal University, Beijing 100875, China. ²Geography and Planning, University at Albany, State University of New York, Albany, NY 12222, USA. ³School of Life Sciences and Global Institute of Sustainability, Arizona State University, Tempe, AZ 85287, USA. ⁴Chinese Research Academy of Environmental Sciences, Beijing 100012, China. ⁵State Key Laboratory of Urban & Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, P. R. China.

Received: 15 July 2014 Accepted: 21 October 2014

Published online: 27 November 2014

References

- Baxter JW, Pickett ST, Dighton J, Carreiro MM (2002) Nitrogen and phosphorus availability in oak forest stands exposed to contrasting anthropogenic impacts. *Soil Biology and Biochemistry* 34:623–633
- Beijing Municipal Bureau of Statistics (2013) Beijing statistical yearbook
- Bremner J, Mulvaney C (1982) Nitrogen—total. In: Miller RH, Keeney DR (eds) *Methods of soil analysis, part 2, chemical and microbiological properties*. American Society of Agronomy, Madison, pp 595–624
- Chen C, Wang EL, Yu Q (2010a) Modeling wheat and maize productivity as affected by climate variation and irrigation supply in North China plain. *Agron J* 102:1037–1049
- Chen X, Xia X, Zhao Y, Zhang P (2010b) Heavy metal concentrations in roadside soils and correlation with urban traffic in Beijing, China. *J Hazard Mater* 181:640–646
- China National Environmental Monitoring Center (1990) The backgrounds value of soil elements in China. China Environment Science, Beijing
- Chon H-T, Ahn J-S, Jung MC (1998) Seasonal variations and chemical forms of heavy metals in soils and dusts from the satellite cities of Seoul, Korea. *Environ Geochem Health* 20:77–86
- Curtin D, Beare MH, Chantigny MH, Greenfield LG (2011) Controls on the extractability of soil organic matter in water over the 20 to 80°C temperature range. *Soil Sci Soc Am J* 75:1423–1430
- Davies R, Hall SJ (2010) Direct and indirect effects of urbanization on soil and plant nutrients in desert ecosystems of the Phoenix metropolitan area, Arizona (USA). *Urban Ecosyst* 13:295–317
- Hamberg L, Malmivaara-Lamsa M, Lehvavirta S, Kotze DJ (2009) The effects of soil fertility on the abundance of rowan (*Sorbus aucuparia* L.) in urban forests. *Plant Ecol* 204:21–32
- Hamzeh MA, Aftabi A, Mirzaee M (2011) Assessing geochemical influence of traffic and other vehicle-related activities on heavy metal contamination in urban soils of Kerman city, using a GIS-based approach. *Environ Geochem Health* 33:577–594
- Harrell MO, Pierce P, Mootr D (1988) Pin oak and silver maple chlorosis treatment with ferric ammonium citrate solution. *J Arboric* 14:156–158
- Huang L, Wang H, Li Y, Lu S (2013) Spatial distribution and risk assessment of phosphorus loss potential in urban-suburban soil of Lishui, China. *Catena* 100:42–49
- Jenerette GD, Wu J, Grimm NB, Hope D (2006) Points, patches, and regions: scaling soil biogeochemical patterns in an urbanized arid ecosystem. *Glob Chang Biol* 12:1532–1544
- Jim CY (1998a) Soil characteristics and management in an urban park in Hong Kong. *Environ Manag* 22:683–695
- Jim CY (1998b) Urban soil characteristics and limitations for landscape planting in Hong Kong. *Landsc Urban Plan* 40:235–249
- Jin J-W, Ye H-C, Xu Y-F, Shen C-Y, Huang Y-F (2011) Spatial and temporal patterns of soil fertility quality and analysis of related factors in urban-rural transition zone of Beijing. *Afr J Biotechnol* 10:10948–10956
- Kaye J, Majumdar A, Gries C, Buyantuyev A, Grimm N, Hope D, Jenerette G, Zhu W, Baker L (2008) Hierarchical Bayesian scaling of soil properties across urban, agricultural, and desert ecosystems. *Ecol Appl* 18:132–145
- Kuang W (2012) Spatio-temporal patterns of intra-urban land use change in Beijing, China between 1984 and 2008. *Chin Geogr Sci* 22:210–220
- Li Z-G, Zhang G-S, Liu Y, Wan K-Y, Zhang R-H, Chen F (2013) Soil nutrient assessment for urban ecosystems in Hubei, China. *Plos One* 8:e75856
- Liu S, Ma K, Fu BJ, Kang Y, Zhang J, Yuxin Z (2003) The relationship between land form, soil characteristic and plant community structure in the Donglingshan mountain region, Beijing. *Acta Phytocologica Sinica* 27:496–502
- Lovell ST, Taylor JR (2013) Supplying urban ecosystem services through multifunctional green infrastructure in the United States. *Landsc Ecol* 28:1447–1463
- Lu R (2000) Analytical methods of soil agrochemistry. China Agricultural Science and Technology, Beijing, pp 85–96
- Lu SG, Bai SQ (2010) Contamination and potential mobility assessment of heavy metals in urban soils of Hangzhou, China: relationship with different land uses. *Environmental Earth Sciences* 60:1481–1490
- Mao Q, Ganlin H, Keming M, Sun Z (2014) Variations of Soil Lead in Different Land Uses Along the Urbanization Gradient in the Beijing Metropolitan Area. *Int J Environ Res Public Health* 11:3199–3214
- McLean E (1982) Soil pH and lime requirement. *Methods of soil analysis. Part 2, Chemical and microbiological properties*. American Society of Agronomy Madison, Wisconsin, USA, pp 199–224
- Messenger S (1986) Alkaline runoff, soil pH and white oak manganese deficiency. *Tree Physiol* 2:317–325
- Moussa AS, van Rensburg L, Kellner K, Bationo A (2009) Exploring differences of soil quality as related to management in semiarid rangelands in the western Bophirima District, North West province, South Africa. *African Journal of Range and Forage Science* 26:27–36
- Neil K, Wu J, Bang C, Faeth S (2014) Urbanization affects plant flowering phenology and pollinator community: effects of water availability and land cover. *Ecol Process* 3:17
- Park S-J, Cheng Z, Yang H, Morris EE, Sutherland M, Gardener BBM, Grewal PS (2010) Differences in soil chemical properties with distance to roads and age of development in urban areas. *Urban Ecosyst* 13:483–497
- Pouyat R, Groffman P, Yesilonis I, Hernandez L (2002) Soil carbon pools and fluxes in urban ecosystems. *Environmental Pollution* 116 Supplement 1:S107–S118
- Pouyat RV, Yesilonis ID, Russell-Anelli J, Neerchal NK (2007) Soil chemical and physical properties that differentiate urban land-use and cover types. *Soil Sci Soc Am J* 71:1010–1019
- Pouyat R, Yesilonis I, Szlavecz K, Csuzdi C, Hornung E, Korsós Z, Russell-Anelli J, Giorgio V (2008) Response of forest soil properties to urbanization gradients in three metropolitan areas. *Landsc Ecol* 23:1187–1203
- Pouyat RV, Szlavecz K, Yesilonis ID, Groffman PM, Schwarz K (2010) Chemical, physical, and biological characteristics of urban soils. In: Aitkenhead-Peterson J, Volder A (eds) *Urban ecosystem ecology*. American Society of Agronomy, Crop Science Society of America, Soil Science Society of America, Madison, pp 119–152
- Raciti SM, Groffman P, Fahey T (2008) Nitrogen retention in urban lawns and forests. *Ecological Applications* 18:1615–1626
- Sauvé S, Hendershot W, Allen HE (2000) Solid-solution partitioning of metals in contaminated soils: dependence on pH, total metal burden, and organic matter. *Environment Science and Technology* 34:1135–1131
- Schwarz K, Pickett STA, Lathrop RG, Weathers KC, Pouyat RV, Cadenasso ML (2012) The effects of the urban built environment on the spatial distribution of lead in residential soils. *Environ Pollut* 163:32–39
- SEPA China (State Environmental Protection Administration of China) (1995) GB15618-1995, Soil Environmental Quality Standards in China (In Chinese). Available at http://kjs.mep.gov.cn/hjbhbz/bzwb/trhj/trhjzlbz/199603/t19960301_82028.htm. Accessed on June 24, 2014
- Setälä H, Bardgett R, Birkhofer K, Brady M, Byrne L, de Ruiter P, de Vries F, Gardi C, Hedlund K, Hemerik L, Hotes S, Liiri M, Mortimer SR, Pavao-Zuckerman M, Pouyat R, Tsiafouli M, van der Putten WH (2013) Urban and agricultural soils: conflicts and trade-offs in the optimization of ecosystem services. *Urban Ecosyst* 17:239–253
- Shang LY, Sun RH, Wang ZM, Ji YH, Chen LD (2012) Assessment of heavy metal pollution in surface sediments of rivers in northern area of Haihe River Basin, China. *Huan Jing Ke Xue* 33:606–611
- Simard R (1993) Ammonium acetate-extractable elements. *Soil sampling and methods of analysis*. Lewis, Boca Raton, pp 39–42
- Su B, Han X, Qu C, Huang J (2001) Net nitrogen mineralization and trification in one pure pine (*Pinus tabulaeformis*) forests and one pine-oak mixed forests in Dongling mountain region. *Acta Phytocologica Sinica* 5:203
- Szolnoki Z, Farsang A, Puskás I (2013) Cumulative impacts of human activities on urban garden soils: origin and accumulation of metals. *Environ Pollut* 177:106–115
- Tiessen H, Moir J, Carter M (1993) Characterization of available P by sequential extraction. *Soil Sampling and Methods of Analysis* 7:5–229
- USDA (United States Department of Agriculture) (2001) Soil quality test kit guide. Available at http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/health/assessment/?cid=nrcs142p2_053873. Accessed on June 24, 2014
- USEPA (United States Environmental Protection Agency) (1995) Method 3052: Microwave assisted acid digestion of siliceous and organically based matrices. Test methods for evaluating solid waste. Available at http://www.epa.gov/osw/hazard/testmethods/sw846/online/3_series.htm. Accessed on June 24, 2014
- Wang B, Zhao S, Xia DS, Yu Y, Tian SL, Jia J, Jiang XR (2011) Characteristics of heavy metal elements and their relationship with magnetic properties of river sediment from urban area in Lanzhou. *Huan Jing Ke Xue* 32:1430–1440

- Whitcomb CE (1987) Establishment and maintenance of landscape plants. *Landscape*
- Wu J (2014) Urban ecology and sustainability: the state-of-the-science and future directions. *Landscape Urban Plan* 125:209–221
- Xia X, Chen X, Liu R, Liu H (2011) Heavy metals in urban soils with various types of land use in Beijing, China. *J Hazard Mater* 186:2043–2050
- Yang L, Li Y, Peng K, Wu S (2014) Nutrients and heavy metals in urban soils under different green space types in Anji, China. *Catena* 115:39–46
- Zhao D, Li F, Yang Q, Wang R, Song Y, Tao Y (2013) The influence of different types of urban land use on soil microbial biomass and functional diversity in Beijing, China. *Soil Use Manag* 29:230–239
- Zhu W-X, Hope D, Gries C, Grimm NB (2006) Soil characteristics and the accumulation of inorganic nitrogen in an arid urban ecosystem. *Ecosystems* 9:711–724

doi:10.1186/s13717-014-0023-8

Cite this article as: Mao et al.: Spatial heterogeneity of urban soils: the case of the Beijing metropolitan region, China. *Ecological Processes* 2014 **3**:23.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
