

RESEARCH

Open Access



# Assessment of above- and belowground carbon pools in a semi-arid forest ecosystem of Delhi, India

Archana Meena<sup>1\*</sup> , Ankita Bidalia<sup>2</sup>, M. Hanief<sup>3</sup>, J. Dinakaran<sup>2</sup> and K. S. Rao<sup>2</sup>

## Abstract

**Background:** Assessment of carbon pools in semi-arid forests of India is crucial in order to develop a better action plan for management of such ecosystems under global climate change and rapid urbanization. This study, therefore, aims to assess the above- and belowground carbon storage potential of a semi-arid forest ecosystem of Delhi.

**Methods:** For the study, two forest sites were selected, i.e., north ridge (NRF) and central ridge (CRF). Aboveground tree biomass was estimated by using growing stock volume equations developed by Forest Survey of India and specific wood density. Understory biomass was determined by harvest sampling method. Belowground (root) biomass was determined by using a developed equation. For soil organic carbon (SOC), soil samples were collected at 0–10-cm and 10–20-cm depth and carbon content was estimated.

**Results:** The present study estimated 90.51 Mg ha<sup>-1</sup> biomass and 63.49 Mg C ha<sup>-1</sup> carbon in the semi-arid forest of Delhi, India. The lower diameter classes showed highest tree density, i.e., 240 and 328 individuals ha<sup>-1</sup> (11–20 cm), basal area, i.e., 8.7 (31–40 cm) and 6.08 m<sup>2</sup> ha<sup>-1</sup> (11–20 cm), and biomass, i.e., 24.25 and 23.57 Mg ha<sup>-1</sup> (11–20 cm) in NRF and CRF, respectively. Furthermore, a significant contribution of biomass (7.8 Mg ha<sup>-1</sup>) in DBH class 81–90 cm in NRF suggested the importance of mature trees in biomass and carbon storage. The forests were predominantly occupied by *Prosopis juliflora* (Sw.) DC which also showed the highest contribution to the (approximately 40%) tree biomass. Carbon allocation was maximum in aboveground (40–49%), followed by soil (29.93–37.7%), belowground or root (20–22%), and litter (0.27–0.59%).

**Conclusion:** Our study suggested plant biomass and soils are the potential pools of carbon storage in these forests. Furthermore, carbon storage in tree biomass was found to be mainly influenced by tree density, basal area, and species diversity. Trees belonging to lower DBH classes are the major carbon sinks in these forests. In the study, native trees contributed to the significant amount of carbon stored in their biomass and soils. The estimated data is important in framing forest management plans and strategies aimed at enhancing carbon sequestration potential of semi-arid forest ecosystems of India.

**Keywords:** Semi-arid forest, Carbon pool, Forest management, Species composition, Basal area, Carbon allocation pattern

\* Correspondence: [archanam417@gmail.com](mailto:archanam417@gmail.com)

<sup>1</sup>Department of Botany, University of Rajasthan, Jaipur, Rajasthan 302004, India

Full list of author information is available at the end of the article

## Introduction

Forests play a significant role in the global carbon (C) cycle and they store large quantities of C in vegetation and soil (Pan et al. 2011). For example, forests store 638 Gt of C in their ecosystem, with 238 Gt in biomass, which accounts for 80% of biomass C of terrestrial vegetation (FAO 2005). Arid and semi-arid regions cover approximately 47% of the earth's surface with an area of 6.5 million km<sup>2</sup> and 4357 Mha of global forest cover (Bastin et al. 2017). The combined effect of climate change and increase in human population will result in severe droughts, desertification, and land degradations in these ecosystems (Huang et al. 2012, 2016). This will further affect the productivity, biodiversity, soil fertility, and organic matter composition which in turn reduces their C storage and sequestration potential. Among the different abiotic factors, soil water availability is considered as a significant variable controlling the soil C storage in arid and semi-arid ecosystems (Meza et al. 2018). Management strategies focusing on the C sequestration by restoring the existing vegetation would be effective in mitigating climate change and its effects on these ecosystems (Malagnoux et al. 2007). To date, only a few studies on biomass and C pools have been carried out in these ecosystems (Bonino 2006; Wagner et al. 2015).

According to good practice guidance developed by the Intergovernmental Panel on Climate Change (IPCC), the total C stock in any forest ecosystem is derived from aboveground biomass (AGB), belowground biomass (BGB) or roots, forest floor litter biomass (LB) or detritus pool, wood debris, and soil organic matter (SOM). Therefore, assessment of C stored in these different pools is crucial to develop a new conservation policy related to C sequestration and combat climate change. AGB and BGB are the living pools of C in forests, which contributes a significant amount of C to the terrestrial ecosystem (Eggleston et al. 2006). The LB contributes only a small fraction of C to the terrestrial pool and hence considered as a minor pool (Ravindranath and Ostwald 2008). The largest and very important C pool in the terrestrial ecosystem is SOM, which contains soil organic carbon (SOC) and plays a great role in the cycling of nutrients and C between the lithosphere and atmosphere (Lal 2005). C pools in various forest ecosystems are strongly influenced by temperature, rainfall, topography (Vayreda et al. 2012), forest type and structure (Wei et al. 2013), tree species composition (Hu et al. 2015), species diversity (Arasa-Gisbert et al. 2018), land use changes, and human-induced disturbances (Canadell et al. 2007). An estimation of the existing pools of C in different forest types is hence required in order to make necessary management strategies related to C sequestration and storage (Johnson and Kern 2002).

Around 3.2 million km<sup>2</sup>, i.e., 12% of the geographical area, is covered by arid zones of India. In India, biomass and C stock has been reported for various ecosystems based on growing stock volume (GSV) data of forest inventories and appropriate conversion factor related to both biomass and C (Ravindranath et al. 1997; Lal and Singh 2000; Chhabra et al. 2002; Manhas et al. 2006; Sharma et al. 2010; Chaturvedi et al. 2011; Dar and Sundarapandian 2015; Salunkhe et al. 2018). However, most of these studies are limited to mainly tropical and temperate ecosystems of India and a very little information is available from semi-arid forests.

Moreover, the urban ecosystem in urban areas plays a dynamic role in reducing air pollution and sequestration of atmospheric carbon dioxide (CO<sub>2</sub>) in their vegetation biomass and soils. Delhi, considered as one of the most polluted cities in the world, has a unique forest ecosystem located on ridges. The ridge areas are the extensions of Aravalli hills in Delhi with a length of 32 km and serve various ecological, environmental, and social functions. Delhi ridge has been notified as reserved forest, managed mainly with the objectives of increasing the forest cover, biodiversity, conservation by public participation, reduction in monoculture plantations, and encroachments. Both north and central ridge forests alone occupied almost 80% of the total forest area of Delhi. The growing urbanization has led to complete loss of the vegetation and only a few areas of the forests have been protected in Delhi. The regeneration of vegetation is further prevented due to excessive grazing and encroachments. Since it is very difficult to increase forest cover in such developed areas, the C sequestration potential of these forests can be enhanced by implementing some management strategies focusing on conservation of forests. However, there is a lack of information on biomass and C pools of this unique forest ecosystem in Delhi. Thus, the main aim of this study was to assess the above- and belowground pools of C in a semi-arid forest ecosystem of Delhi. The present study has the following objectives: (1) to estimate the C stock in different components (C pools) of a semi-arid forest ecosystem of Delhi, (2) to estimate the C allocation pattern in different pools, and (3) to analyze various factors influencing the C storage pattern in these components.

## Methods

### Study area

The state of Delhi, comprising National Capital Territory (NCT), lies between 28°24'17" N and 28° 53' 00" N latitudes and 76° 50' 24" E and 77° 24' 17" E longitudes and covers an area of 1483 km<sup>2</sup>. Physiogeographically, the area is dominated by river Yamuna, the Aravalli range, and the plains in between, formed by alluvium deposits of

recent origin. About 25% of the total area of NCT is in rural and the remaining 75% is urban. As per Census India 2011, the population of Delhi is 16.8 million with a decadal growth of 21.21%. As per the Land Use Statistics, Ministry of Agriculture, Government of India (2013–2014), net sown area, current fallow, and culturable wasteland is 221.4, 119, and 98.9 km<sup>2</sup> covering 15%, 8.07%, and 6.71% of its total geographical area, respectively. Forest's land use covers 14.8 km<sup>2</sup> with 1% of the total geographical area. A major chunk of land, i.e., 927 km<sup>2</sup> representing 62.85%, is not available for agriculture. With rapid urbanization in Delhi, there is a continuous decrease in rural area and agricultural land.

Delhi ridge which is the extension of Aravalli hills is estimated to be 2.4 billion years old. The ridge recorded a massive afforestation during nineteenth century and has been notified as reserved forests under the Indian Forest Act, 1927. However, due to rapid urbanization, there was a decrease in ridge forest area between 1920 and 1930. The ridge forest was, then again, declared as reserve forest in 1980 in order to limit the anthropogenic activities. At present, the recorded forest area of Delhi state is 102 km<sup>2</sup>, which constitute 6.88% of its geographical area, with reserve and protected forests comprising 76.48% and 23.52% of total forest area, respectively (FSI 2017). The forest type is tropical thorny forest (Champion and Seth 1968). The vegetation is mainly dominated by middle storied thorny trees with open patches having scattered distribution. The soil type on the ridge has been reported as sandy loam to loam (Chibbar 1985). *Prosopis juliflora* (Sw.) DC, an exotic species, is the dominant tree in the forest. *Acacia nilotica* (L.) Delile, *Acacia leucophloea* (Roxb.) Willd., *Salvadora oleoides* Decne, and *Cassia fistula* L. are among the commonly found native trees (Sinha 2014; Meena et al. 2016). The commonly growing shrubs in the forests are *Justicia adhatoda* L., *Capparis sepiaria* L., *Carissa spinarum* L., *Jatropha gossypifolia* L., and *Opuntia dillenii* L. Two sites were selected for this study, (1) north ridge forest (NRF) and (2) central ridge forest (CRF). The NRF has an area of 87 ha and situated at 28° 36' latitudes and 77° 41' E longitudes. The CRF has an area of 864 ha and situated at 28° 41' latitudes and 77° 12' E longitudes. The climate of the study area is semi-arid and characterized by hot-dry summers (April to June), monsoon (July to October) and cool-dry winters (November to January). The study area received a total annual precipitation of 720 mm with 31 and 19 °C as mean maximum and minimum air temperature during 2012 (IMD, 2012). About 87% of the total rain was received during the monsoon. The soil in the study sites are dry and sandy in nature.

#### Phytosociology of the study sites

A total of 15 and 25 plots of 10 × 10 m were laid randomly in NRF and CRF sites respectively. The

phytosociological data of the study sites were collected during August and September in the year 2012. All the trees with the diameter ≥ 10 cm DBH (1.37 m above from the base) occurring in each plot were measured and identified to the species level. All the trees were considered as individuals as per Knight (1975) and individuals with DBH < 10 cm were recorded as seedling/saplings (Pande et al. 1988). The phytosociological parameters were evaluated using the standard methods suggested by Misra (1968). Tree density (TD) was estimated by dividing the total number of individuals of a species with a total number of quadrats studied. The basal area (BA) of each tree was calculated as the ratio of CBH<sup>2</sup> (squared circumference at breast height) to 4π.

#### Estimation of tree biomass and C stock

To determine the aboveground tree biomass (AGTB), the GSV (m<sup>3</sup> ha<sup>-1</sup>) of each species was first estimated by using volume tables or equations (Table 1), determined by Forest Survey of India (FSI 1996). These equations were developed using multiple regression methods considering the DBH along with tree height or form factor. For the trees where volume equations were not available, the general or local area-based equations were used. The estimated GSV of the tree was then converted to AGTB (Mg ha<sup>-1</sup>) by multiplying GSV with specific wood density (g cm<sup>-3</sup>) of the respective species (Rajput et al. 1996). Global wood density database was used for species-specific wood density values (Chave et al. 2009; Zanne et al. 2009). The BGB (fine and coarse roots) was estimated using regression equations suggested by Cairns et al. (1997) as:

$$\text{BGB (Mg ha}^{-1}\text{)} = \exp \{-1.059 + 0.884 \times \ln (\text{AGB}) + 0.284\}$$

The total tree biomass (TB) (Mg ha<sup>-1</sup>) is described as the sum of AGB and BGB.

#### Understory biomass

The understory biomass, i.e., shrubs (woody species other than trees with less than 1 m height) and herbs, was estimated by randomly laying 5 × 5 m and 1 × 1 m quadrat for shrubs and herbs, respectively. The shrub biomass (SB) was estimated by harvesting method, where 10% of each species of shrub was harvested and fresh weight of the harvested sample was measured immediately with an electronic balance in the field. For the herbaceous biomass (HB), all the herbaceous vegetation falling in 1 × 1 m quadrat was harvested and fresh weight was measured immediately in the field. The representative samples of both herbs and shrubs were taken to the laboratory, where they were oven dried at 65 °C for 48 h. The dry weight of the sample was then estimated.

**Table 1** Volume equation and wood density of tree species

Tree species	Volume equation*	Wood density (g cm <sup>-3</sup> )**
<i>Acacia leucophloea</i> (Roxb.) Willd.	$\text{sqrt } V = -0.00142 + 2.61911D - 0.54703 \times \text{sqrt } D$	0.9
<i>A. modesta</i> Wall.	$\text{sqrt } V = -0.00142 + 2.61911D - 0.54703 \times \text{sqrt } D$	0.9
<i>A. nilotica</i> (L.) Delile	$\text{sqrt } V = -0.00142 + 2.61911D - 0.54703 \times \text{sqrt } D$	0.9
<i>Albizia lebbeck</i> (L.) Bent.	$V = 0.00471 + 1.79326D^2$	0.53
<i>Azadirachta indica</i> Juss.	$V = 0.00471 + 1.79326D^2$	0.7
<i>Bauhinia purpurea</i> L.	$V = 0.00471 + 1.79326D^2$	0.67
<i>Butea monosperma</i> (Lam.) Taub.	$V = 0.00471 + 1.79326D^2$	0.48
<i>Cassia fistula</i> L.	$V = 0.066 + 0.287D^2$	0.64
<i>Cordia dichotoma</i> G. Forst.	$V = 0.00471 + 1.79326D^2$	0.53
<i>Crateva religiosa</i> Forst.f.	$V = 0.00471 + 1.79326D^2$	0.53
<i>Ficus drupacea</i> Thunb.	$\text{sqrt } V = 0.03629 + 3.95389 \times D - 0.84421\text{sqrt } D$	0.39
<i>F. racemosa</i> L.	$\text{sqrt } V = 0.03629 + 3.95389 \times D - 0.84421\text{sqrt } D$	0.39
<i>Holoptelea integrifolia</i> Planch.	$V = 0.00471 + 1.79326D^2$	0.64
<i>Pithecellobium dulce</i> (Roxb.) Benth.	$V = 0.00471 + 1.79326D^2$	0.5
<i>Pongamia pinnata</i> (L.) Pierre	$V = 0.00471 + 1.79326D^2$	0.82
<i>Prosopis juliflora</i> (Sw.) DC	$V = 0.00471 + 1.79326D^2$	0.73
<i>Salvadora oleoides</i> Decne.	$V = 0.00471 + 1.79326D^2$	0.59
<i>Syzygium cumini</i> (L.) Skeels.	$V = 0.00471 + 1.79326D^2$	0.69

\*(FSI 1996)

\*\*(Zanne et al. 2009)

**Litter biomass (LB)**

The forest LB was estimated by collecting the litter at quarterly interval using specially designed plastic trays, laid randomly within 10 × 10 m quadrat and pooled for further analysis. The fresh weight of the litter (leaves and branches) was taken in the field. The representative samples were brought to the laboratory where the samples were oven dried at 65 °C for 48 h and subsequently the dry weight was measured.

**Soil organic carbon (SOC)**

Soil samples were collected during the same period from each plot at two depths, i.e., 0–10 and 10–20 cm. The soil samples were collected randomly from three points from each plot and a composite was made. For SOC analysis, the soil samples were dried at 50 °C for 24 h and sieved through a 2-mm sieve. The sieved samples were further ground in a mortar with pestle and then analyzed using CHNS Analyzer (Elementar vario).

For bulk density, undisturbed soil samples were collected with a soil corer of known volume (31.4 cm<sup>3</sup>) which was inserted at 0–10 and 10–20-cm depth. The soil samples were oven dried at 105 °C for 72 h and the dry weight was measured. The coarse rock fragments if present in the soil were separated and weighed. SOC stock was calculated for each layer from 0 to 10 cm and 10–20-cm depth based on the bulk density and SOC concentration by using the following equation:

$$\text{SOC stock (Mg C ha}^{-1}\text{)} = K_d \times \text{BD} \times \text{SOC}(\%) \times \text{CF} \times 10$$

where  $K_d$  is the depth of soil (cm), BD is the bulk density (g cm<sup>-3</sup>), SOC (%) is soil organic carbon, and CF is a correction factor for coarse fragments (> 2-mm particles) (Borah et al. 2015).

**Estimation of C stock**

The C stock of the tree species was determined as:

$$\text{Carbon (Mg C ha}^{-1}\text{)} = \text{Biomass (Mg ha}^{-1}\text{)} \times C\%$$

where  $C$  is the carbon concentration of the corresponding vegetation. Since it was difficult to separate the different components of the tree for  $C$  estimation, a universal coefficient of 0.475 was used for tree  $C$  estimation (Raghubanshi 1991; Singh and Chand 2012), indicating approximately 47.5% of  $C$  in dry plant biomass (Westlake 1963). The understory (shrubs and herbs) and forest LB was estimated to be 50% of the biomass (Dar and Sundarapandian 2015).

The total ecosystem  $C$  was taken as the sum of  $C$  content in all the pools, i.e., AGB, BGB, SB, HB, LB, and SOC.

**Statistical analysis**

All the statistical analyses were done using the SPSS software package (SPSS version 16, SPSS Inc., Chicago 1 L, USA). The independent  $t$  tests were performed to test

the significant differences in TD, BA, AGTB, BGB, SB, HB, and C content between NRF and CRF. For SOC, the significance was tested between the two forest sites and at different depths among a forest. The significance was tested at  $p < 0.05$ .


**Results**

**Forest’s tree stand and biomass**


TD and BA in different DBH classes are shown in Fig. 1a and b, respectively. While the total TD was high in the CRF (684 individuals ha<sup>-1</sup>) than NRF (633 individuals ha<sup>-1</sup>), most of the trees in CRF are distributed in lower DBH classes. In NRF, the distribution of the trees on the basis of the DBH class was wider with comparatively higher TD in DBH above 60 cm. In both the sites, individuals within diameter class 11–20 cm were dominated with TD of 240 and 328 individuals ha<sup>-1</sup> in NRF and CRF, respectively. The maximum BA was contributed by individuals in DBH class 31–40 cm (8.87 m<sup>2</sup> ha<sup>-1</sup>) in NRF and 11–20 cm (6.08 m<sup>2</sup> ha<sup>-1</sup>) in CRF site. The distribution of TB in different DBH classes is shown in Fig. 2c. The maximum biomass was observed for trees belonging to DBH class 11–20 cm in both NRF (24.25 Mg ha<sup>-1</sup>) and CRF (23.57 Mg ha<sup>-1</sup>). The species-wise contribution to the TB in both forests is given in Table 2. Highest contribution (~ 40%) was observed by *P. juliflora*, having maximum biomass at both sites (44.39 and 28.68 Mg ha<sup>-1</sup> in NRF and CRF, respectively). The other associated tree species also contribute significantly to the total TB like *Acacia leucophloea* (18.31 and 11.55 Mg ha<sup>-1</sup> in NRF and CRF, respectively), *Pongamia pinnata* (L.) Pierre (8.22 and 3.26 Mg ha<sup>-1</sup> in NRF and CRF, respectively), and *Azadirachta indica* Juss. (6.72 and 3.12 Mg ha<sup>-1</sup> in NRF and CRF, respectively).

**Biomass in different components or pools**

The biomass in the different components of the two different study sites is given in Table 3. The AGTB for NRF and CRF was estimated as 75.24 and 44.95 Mg ha<sup>-1</sup> and BGB was 32.61 and 24.25 Mg ha<sup>-1</sup>, respectively (Table 3). The TB, AGTB, and BGB were significantly different between the two forest sites ( $p < 0.05$ ). The understory biomass (shrubs and herbs) was 1.63 and 1.29 Mg ha<sup>-1</sup> in NRF and CRF, with a mean of 1.46 Mg ha<sup>-1</sup>. The SB was estimated to be 1.15 and 0.80 Mg ha<sup>-1</sup> in NRF and CRF, with a mean of 0.98 Mg ha<sup>-1</sup>. The HB ranged from 0.48 to 0.49 Mg ha<sup>-1</sup> in NRF and CRF, with a mean of 0.49 Mg ha<sup>-1</sup>. However, we did not find a significant difference ( $p > 0.05$ ) for the biomass of SB and HB between the two forests. The forest LB was significantly higher in CRF (0.64 Mg ha<sup>-1</sup>) as compared to NRF (0.40 Mg ha<sup>-1</sup>), with the mean of 0.52 Mg ha<sup>-1</sup>.



**Fig. 1** Distribution of tree a density, b basal area, and c biomass in DBH classes. DBH diameter at breast height, NRF north ridge forest, CRF central ridge forest



**Fig. 2** Contribution of carbon in different pools. AGB aboveground biomass, BGB belowground biomass, LB litter biomass, SOC soil organic carbon

**Table 2** Biomass and carbon content in different tree species

Plant species	AGTB (Mg ha <sup>-1</sup> )		BGB (Mg ha <sup>-1</sup> )		TC (Mg ha <sup>-1</sup> )	
	NRF	CRF	NRF	CRF	NRF	CRF
<i>Acacia leucophloea</i> (Roxb.) Willd.	13.03	8.03	5.28	3.52	8.7	5.49
<i>A. modesta</i> Wall.	0.17	–	0.10	–	0.13	–
<i>A. nilotica</i> (L.) Delile	–	10.25	–	4.58	–	7.04
<i>Albizia lebbeck</i> (L.) Bent.	2.29	–	0.96	–	1.54	–
<i>Azadirachta indica</i> Juss.	4.59	2.09	2.13	1.03	3.19	1.49
<i>Bauhinia purpurea</i> L.	0.14	–	0.09	–	0.11	–
<i>Butea monosperma</i> (Lam.) Taub.	0.12	0.61	0.07	0.36	0.09	0.46
<i>Cassia fistula</i> L.	0.29	0.03	0.18	0.02	0.22	0.02
<i>Cordia dichotoma</i> G. Forst.	0.54	–	0.28	–	0.39	–
<i>Crateva religiosa</i> Forst.f.	0.15	–	0.09	–	0.12	–
<i>Ficus drupacea</i> Thunb.	–	3.07	–	1.34	–	2.09
<i>F. racemosa</i> L.	0.90	–	0.46	–	0.65	–
<i>Holoptelea integrifolia</i> Planch.	10	–	3.83	–	5.14	–
<i>Pithecellobium dulce</i> (Roxb.) Benth.	0.21	–	0.12	–	0.16	–
<i>Pongamia pinnata</i> (L.) Pierre	5.70	2.12	2.52	1.14	3.9	1.55
<i>Prosopis juliflora</i> (Sw.) DC	30.22	14.17	13.72	9.98	21.09	13.62
<i>Salvadora oleoides</i> Decne.	0.66	4.58	0.36	2.28	0.49	3.26
<i>Syzygium cumini</i> (L.) Skeels.	6.23	–	2.42	–	4.11	–

NRF north ridge forest, CRF central ridge forest, AGTB aboveground tree biomass, BGB belowground biomass, TC tree carbon

### Allocation pattern of C in different pools

The C content in different compartments (pools) in both forests is given in Table 4 and the pattern of allocation is shown in Fig. 2. Maximum C allocation was observed in AGB pool (sum of AGTB, SB, and HB) contributing 49.02 and 40.54% in NRF and CRF, respectively, to the total forest C stock. This was followed by SOC pool, with an allocation of 29.93 and 37.7% in NRF and CRF, respectively. BGB contributed 20.77 and 21.26% of C in NRF and CRF, respectively, followed by LB with an allocation of 0.27 and 0.59% of C in NRF and CRF, respectively. Within the selected forest sites, the C storage was found to be high in AGB (36.56 Mg C ha<sup>-1</sup>), BGB (15.49 Mg C ha<sup>-1</sup>), and SOC (74.57 Mg C ha<sup>-1</sup>), pools in NRF than CRF, whereas the C

storage in the LB was high in CRF (0.32 Mg C ha<sup>-1</sup>) as compared to NRF (0.2 Mg C ha<sup>-1</sup>).

SOC was found to be high in upper (0–10 cm) as compared to lower soil layers (10–20 cm) in both forest sites (Table 5). The SOC stock at 0–10-cm depth was estimated to be 26.60 and 24.36 Mg C ha<sup>-1</sup> in NRF and CRF sites, respectively. The SOC stock at 10–20-cm depth was estimated to be 18.04 and 16.50 Mg C ha<sup>-1</sup> in NRF and CRF sites, respectively. The mean SOC stock from 0 to 20-cm depth was estimated to be 22.32 and 20.43 Mg C ha<sup>-1</sup> for NRF and CRF sites, respectively. The SOC stock both in

**Table 3** Biomass in different ecosystem components

Ecosystem components	Biomass (Mg ha <sup>-1</sup> )	
	NRF	CRF
TB	107.85 ± 2.68 a	69.2 ± 2.13 b
AGTB	75.24 ± 2.08 a	44.95 ± 1.84 b
BGB	32.61 ± 0.63 a	24.25 ± 0.6 b
SB	1.15 ± 0.52 a	0.80 ± 0.3a
HB	0.48 ± 0.03 a	0.49 ± 0.06 a
LB	0.40 ± 0.06 a	0.64 ± 0.07 b

Each value represents mean ± SE. Letters denotes significant difference at  $p < 0.05$ . NRF north ridge forest, CRF central ridge forest, AGTB aboveground tree biomass, BGB belowground biomass, SB shrub biomass, HB herb biomass, LB litter biomass

**Table 4** Carbon content in different pools of forests

Forest carbon pools	Carbon content (Mg C ha <sup>-1</sup> )		Mean (Mg C ha <sup>-1</sup> )
	NRF	CRF	
AGTB	35.74 ± 0.98 a	21.35 ± 0.76 b	28.55
SB	0.57 ± 0.26 a	0.4 ± 0.15 a	0.52
HB	0.25 ± 0.02 a	0.26 ± 0.03 a	0.26
AGB	36.56	22.01	29.28
BGB	15.49 ± 0.30 a	11.52 ± 0.25 b	12.59
LB	0.2 ± 0.03 a	0.32 ± 0.04 b	0.26
SOC	22.32 ± 3.26 a	20.43 ± 1.58 a	21.36
Total carbon stock	74.57	54.28	63.49

Each value represents mean ± SE. Letters denotes significant difference at  $p < 0.05$ . NRF north ridge forest, CRF central ridge forest, AGTB aboveground tree biomass, BGB belowground biomass, SB shrub biomass, HB herb biomass, LB litter biomass, SOC soil organic carbon

0–10 and 10–20 cm depth did not vary significantly ( $p > 0.05$ ) between the two forest sites. Furthermore, no significant differences ( $p > 0.05$ ) were observed in SOC stock between two depths in NRF forest sites. However, in CRF site, the SOC stock between two depths varied significantly ( $p < 0.05$ ).

## Discussion

The estimated total biomass of the forest sites range between 71.13 and 109.88 Mg ha<sup>-1</sup>, with an average of 90.51 Mg ha<sup>-1</sup>, within the range reported for Indian forest systems (27.4 to 251.8 Mg ha<sup>-1</sup>) (Chhabra et al. 2002), but is higher than the average (40 Mg ha<sup>-1</sup>) value reported for thorn forest system (Ravindranath et al. 1997). This clearly shows that the studied forest system in Delhi has the potential to store more C in its plant biomass. The C stored in different pools was estimated as aboveground (29.28 Mg C ha<sup>-1</sup>), roots or belowground (12.59 Mg C ha<sup>-1</sup>), litter (0.26 Mg C ha<sup>-1</sup>), and SOC (21.36 Mg C ha<sup>-1</sup>). The estimated average forest standing biomass C (sum of AGB and BGB), i.e., 41.87 Mg C ha<sup>-1</sup>, was lower than the values reported for tropical dry forests of Asia (Gibbs et al. 2007; IPCC 2006) and forests from Mexico (Dai et al. 2014) and Ethiopia (Solomon et al. 2017, 2018), but higher than forests from Africa (Gibbs et al. 2007; IPCC 2006) and Brazil (Júnior et al. 2016). Carbon in AGB was also lower than that of a semi-arid (*Picea crassifolia*) forest in the North Eastern Tibet (Wagner et al. 2015). The mean standing biomass C for the present forests is within the range reported for tropical dry deciduous (12.79–62.48 Mg C ha<sup>-1</sup>) and evergreen forests (18.85–48.58 Mg C ha<sup>-1</sup>) but higher than tropical thorn forests (4.91–13.3 Mg C ha<sup>-1</sup>) from India (FSI 2017) (Table 6).

Tree species composition also affects the total biomass and C stock of forest (Yang et al. 2005; Borah et al. 2015; Solomon et al. 2017). In the present study, *P. juliflora* was found to be the most dominant tree species in both forest sites, having the highest TD, BA (Meena et al. 2016),

biomass, and C stock. However, the other associated tree species like *Holoptelea integrifolia*, *Acacia leucophloea*, *Acacia nilotica*, *Azadirachta indica*, *Syzygium cumini*, *Pongamia pinnata*, and *Albizia lebbeck*, though having a low TD, contribute significantly to the total biomass in NRF site. High tree species diversity in NRF than CRF further indicates its effect on forest productivity, increased biomass, and C stock. Management interventions and increased soil moisture content in NRF than in CRF promote the establishment of different species (Meena et al. 2016). Additionally, differences in TD may also contribute to differences in C density among the forest types (Baker et al. 2004; Behera et al. 2017). BGB/AGB ratio indicates the biomass allocation and stability. A high ratio was observed for species like *Bauhinia purpurea* (0.64), *Cassia fistula* (0.62), *Acacia modesta* (0.58), and *Butea monosperma* (0.62) in these forests. Furthermore, a higher ratio for *P. juliflora* in CRF (0.7) than in NRF (0.45) suggested a deep rooting system allowing more penetration and establishment in dry areas. The low ratio in NRF (0.43) than in CRF (0.54) indicated trees with a shallow rooting system, which easily uprooted by the wind. Furthermore, the higher investment to BGB in CRF than in NRF could be due to lower water tables in CRF.

The significant role of understory pool (shrubs and herbs) in C sequestration and maintaining the biodiversity of the forest necessitates its estimation in C stock studies (Hou et al. 2015; Yue et al. 2018). The contribution of understory (shrubs and herbs) C storage to total forest C ranges between 1.09–2.02%, which is within the values (3%) as reported by Brown et al. (1997). However, the allocation of understory biomass is high when compared to the other forests of India, i.e., 0.13% in temperate forests of Kashmir (Dar and Sundarapandian 2015), 0.15% in the humid tropics in the Northeast (Baishya and Barik 2011), 0.1–0.3% in spruce forests in Northwest China (Yue et al. 2018), and also with similar forest type of *Picea crassifolia* forests (0.4%) in the Qilian Mountains (northeastern Tibetan Plateau) (Wagner et al. 2015). The difference in the understory pool of the arid zone could be attributed to species composition, canopy cover, nutrients, and light (Abdallah and Chaieb 2012). As the studied forest is an open forest covering 8.09% of the total forest cover (11.88%), the lower TD favors the growth of ground vegetation, which subsequently increases the biomass and C storage in this pool.

Forest floor litter represents the detritus C considered as the most active pool in the forest C cycle (Yanai et al. 2003). The contribution of C in LB pool to the total C stock is smaller (0.2–1%) as compared to other components but is quite significant (FAO 2005). The estimated litter C is within the range reported for the tropical thorn forest of India (FSI 2017). The contribution of

**Table 5** Soil organic carbon content at different depths

Parameters	Soil depth (cm)	Sites	
		NRF	CRF
C (g kg <sup>-1</sup> )	0–10	16.83 ± 2.84	15.20 ± 2.03
	10–20	10.93 ± 1.75	9.19 ± 1.17
BD (g cm <sup>-3</sup> )	0–10	1.58	1.56
	10–20	1.65	1.62
SOC stock (Mg ha <sup>-1</sup> )	0–10	26.6 ± 4.49* a	24.36 ± 2.09*a
	10–20	18.04 ± 2.88* a	16.5 ± 1.49*b

Each value represents mean ± SE. Letters in parenthesis denotes significant difference between two depth and \* denotes significant difference between forest sites at  $p < 0.05$ . NRF north ridge forest, CRF central ridge forest, SOC soil organic carbon, BD bulk density

**Table 6** Comparison of carbon content ( $\text{Mg C ha}^{-1}$ ) in standing biomass, litter, and soil of dry forests

Forest type/region	Standing biomass	LB	SOC	Source
Tropical dry forest of Asia	120; 78–96			Gibbs et al. (2007); IPCC (2006)
Tropical dry forests of Africa	17			Gibbs et al. (2007)
Tropical dry forest, Brazil	19.27	2.62		Júnior et al. (2016)
Dry forest Tigray, Ethiopia	58.11			Solomon et al. (2017)
Dry Afromontane forest in Northern Ethiopia	15.59–77.19	1.68–2.25	87.55–102.33	Solomon et al. (2018)
Secondary tropical dry forest in the Yucatan Peninsula, Mexico	56.6			Dai et al. (2014)
Semi-arid ( <i>Picea crassifolia</i> ) forest in northeastern Tibet	55	3	306 (0–100 cm)	Wagner et al. (2015)
Tropical dry forest of Javadi Hills, India	52–116			Naveenkumar et al. (2017)
Tropical dry deciduous forest, India	93.8			Singh (1990)
Tropical thorn forest, India	40			Salunkhe et al. (2018)
Tropical dry forests of Eastern Ghats, India	6.98–257.25			Sahu et al. (2016)
Tropical dry deciduous forest, Eastern Ghats, India			16.92–44.65	Gandhi and Sundarapandian (2017)
Tropical dry deciduous forests in Central India	48.97–214.97			Joshi and Dhyani (2018)
Tropical dry deciduous forest, India	12.79–62.48	0.42–6.49	30–59.23	FSI (2017)
Tropical dry evergreen forest, India	18.85–48.58	0.9–1.91	35.08–89.01	FSI (2017)
Tropical thorny forest, India	4.91–13.30	0.76–2.18	19.43–30.17	FSI (2017)
Semi-arid forest in Delhi, India	41.87	0.26	21.36 (0–20 cm)	Present study

LB litter biomass, SOC soil organic carbon

forest litter pool was similar or close to the values reported for different forest types by Baishya and Barik (2011) (0.5%); Zhang et al. (2013) (2–4%); Dar and Sundarapandian (2015) (2.06%), and from a similar forest type by Wagner et al. (2015) (0.9%). The amount of litter C is controlled by various factors like age and density of trees (Yue et al. 2018), soil nutrient levels (Ovington 1956), species composition, quantity and quality of annual litter input (Zhang and Wang 2010), decomposition rate (Chaturvedi and Singh 1987; Taylor et al. 2007), anthropogenic disturbances, and management history.

The average SOC stock for the forest up to 20-cm depth was estimated as  $21.36 \text{ Mg ha}^{-1}$ . These values were lower than those reported for tropical dry deciduous forests ( $37.5 \text{ Mg ha}^{-1}$  for 50 cm, Chhabra et al. 2003), southern thorn forest of India ( $76.85 \text{ Mg ha}^{-1}$ , Ramachandran et al. 2007), forests of Kolli hills of Tamil Nadu ( $175\text{--}369 \text{ Mg ha}^{-1}$ , Mohanraj et al. 2011), tropical dry forest of Brazil ( $47.73\text{--}61.6 \text{ Mg ha}^{-1}$ , Santos et al. 2016), and a semi-arid thicket of South Africa ( $168 \text{ Mg ha}^{-1}$ , Mills et al. 2005). However, these estimated SOC values are comparable to those reported for other tropical dry forests of India at 0–30-cm depth for Sathnur reserve forest ( $16.92\text{--}44.65 \text{ Mg ha}^{-1}$ , Gandhi and Sundarapandian 2017), Uttar Pradesh ( $21.8 \text{ Mg ha}^{-1}$ , Chaturvedi et al. 2011), thorn forest of India ( $19.43\text{--}30.17 \text{ Mg ha}^{-1}$ , FSI 2017), and also with tropical dry forest of Australia ( $29.98 \text{ Mg ha}^{-1}$ , Gray et al. 2015). The amount of SOC content is determined by the rate of organic matter inputs, accumulation, rate of mineralization in different organic C pools

(Post and Kwon 2000), stand type, and stand age (Cao et al. 2018). The overall contribution of SOC to total forest C stock was found to be 29–38%. The reduced SOC concentrations in the present forest type probably result from the high lignin-containing litter from *P. juliflora* and other associated semi-arid trees and low soil moisture content which limits the soil microbial activity and decomposition of organic matter (Wagner et al. 2015; Yue et al. 2018). The rate of litter decomposition also significantly control the SOC stock in a forest. Exposure to high solar radiations in the semi-arid ecosystems results in loss of litter mass and C via altering the rate of litter decomposition (Zhang et al. 2008). UV exposure along with interannual variability in precipitation changes the foliar traits and litter chemistry by causing photo-degradation of lignin and C, hence affecting the overall litter decomposition in these ecosystems (Gaxiola and Armesto 2015).

Over the years, the studied forests are influenced by anthropogenic disturbances including encroachments, resource exploitation, construction, deforestation, over-grazing, exotic plantation, and urbanization which resulted in forest fragmentation and have altered the forest structure and species composition (Sinha 2014). This has serious impacts on future biomass and C storage potential of these forests (Pan et al. 2011). A similar study by Muhati et al. (2018) in a sub-humid disturbed montane forest of Kenya reported decreased values of AGB than indigenous dry tropical forests, which was suggested due

to the dominance of young trees having low BA and height, a characteristic of a regenerating forest. In our study, lower values of C in AGB could be attributed to a high TD of individuals belonging to lower DBH classes. However, the presence of more trees in higher DBH classes has significantly increased the biomass in NRF as compared to CRF, indicating the importance of mature trees in C storage in a forest. The contribution of large trees (DBH > 70 cm) to AGB in the forest was reported up to 50% in previous studies (Brown and Lugo 1992; Brown et al. 1996; Clark and Clark 1996; Baishya et al. 2009). Therefore, the increased volume of mature trees has made a noteworthy contribution to total AGB and C sequestration in NRF site as observed in other studies (Stephenson et al. 2014; Kauppi et al. 2015; Behera et al. 2017). The C stock of tree species correlates positively with BA, TD, diversity, and forest productivity (Baishya et al. 2009; Borah et al. 2015; Joshi and Dhyani 2018). Furthermore, the age of forest stand also influences the biomass and correlates positively with forest C stock in previous studies, indicating an increase in C storage with stand age (Wei et al. 2013; Köhl et al. 2017). The older stand age of these forests results in temporal net primary productivity (NPP) accumulation and increases the overall tree C storage (Chen et al. 2016). Overall, our study shows that the living portions (AGB + BGB) of the mature and young trees in the studied forest ecosystem in Delhi could play a significant role in the storage of more C in their plant biomass than soils. The SOC pool significantly increases the total C stock in such forest types and therefore has the potential for C sequestration. The study also emphasized the minor pools of C in the forest, i.e., the understory C pool and detritus pool which although have a small but significant contribution to the total C stock in the studied forest.

## Conclusion

The study estimated the C content in different pools in a semi-arid forest ecosystem in Delhi, in order to understand their allocation and C sequestration potential. Our study estimated maximum storage of C in plant biomass contributing 40–49% of the total forest C stock suggesting it as a great potential pool of C storage in these forests. Furthermore, it was found that the C storage potential in the forests is influenced by tree basal area, density, and species composition. Maximum biomass contribution by trees in lower DBH classes revealed the importance of young trees as major sinks of C in these forests. However, our results emphasize the protection of old-growth mature tree species having low TD but high BA as these are under the threat of extinction. In the present forest, *P. juliflora*, being the most dominant tree species, adapts better to moisture stress condition and showed highest biomass and C storage. However, attention should be given for the conservation of other native tree species like *Acacia leucophloea*,

*Acacia nilotica*, and *Albizia lebbek* showing high storage of biomass and C. The study also focuses on SOC pool contributing 29–38% to total C stock of forest, which suggests it as a potential sink in C sequestration. The minor pools (LB) also contribute significantly to the total C storage in these forests and therefore should be considered in C stock studies. The studied semi-arid forest is notified as reserved forest, but over the years, forest destruction for construction and expansion leads to the loss of indigenous trees. The C stock studies are difficult in such areas as the forests have more fragmented patches resulting in greater uncertainty and difficulty in estimation of tree biomass. Our study shows the importance of semi-arid areas in C stock and provides necessary data to the researchers and forest managers for developing management plans in arid and semi-arid forests of India.

## Abbreviations

AGB: Aboveground biomass; AGTB: Aboveground tree biomass; BA: Basal area; BGB: Belowground biomass; C: Carbon; CRF: Central ridge forest; DBH: Diameter at breast height; GSV: Growing stock volume; HB: Herbaceous biomass; LB: Litter biomass; NRF: North ridge forest; SB: Shrub biomass; SOC: Soil organic carbon; TB: Tree biomass; TD: Tree density

## Acknowledgments

Authors thankfully acknowledge the Council of Scientific and Industrial Research (CSIR), University Grants Commission (UGC) (UGC-DSK-PDF (BSR)/BL/16-17/0146) and Department of Science and Technology (DST), India for financial support. We also thanks University of Delhi for providing Research and Development for providing grant for doctoral research program.

## Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

## Authors' contributions

AM proposed the idea and conducted the field sampling, data collection, laboratory analysis, data interpretation, and manuscript writing. AB and MH carried out field sampling and data collection. JD helped in the analysis of data and edited the manuscript. KSR guided the study, interpreted the results, and critically reviewed the idea. All authors read and approved the final manuscript.

## Ethics approval and consent to participate

Not applicable.

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Author details

<sup>1</sup>Department of Botany, University of Rajasthan, Jaipur, Rajasthan 302004, India. <sup>2</sup>Natural Resource Management Laboratory, Department of Botany, University of Delhi, North Campus, Delhi 110007, India. <sup>3</sup>Department of Botany, Baba Ghulam Shah Badshah University, Rajouri, J&K 185234, India.

Received: 23 December 2018 Accepted: 27 February 2019

Published online: 04 April 2019

## References

- Abdallah F, Chaieb M (2012) The influence of trees on nutrients, water, light availability and understorey vegetation in an arid environment. *Appl Veg Sci* 15(4):501–512
- Arasa-Gisbert R, Vayreda J, Román-Cuesta RM, Villela SA, Mayorgab R, Retana J (2018) Forest diversity plays a key role in determining the stand carbon stocks of Mexican forests. *For Ecol Manag* 415–416:160–171
- Baishya R, Barik SK (2011) Estimation of tree biomass, carbon pool and net primary production of an old-growth *Pinus kesiya* Royle ex. Gordon forest in north-eastern India. *Ann For Sci* 68(4):727–736
- Baishya R, Barik SK, Upadhya K (2009) Distribution pattern of aboveground biomass in natural and plantation forests of humid tropics in Northeast India. *Trop Ecol* 50(2):295–304
- Baker T, Phillips OL, Malhi Y, Almeida S, Arroyo L, Fiore AD, Killeen TJ, Laurance SG, Laurance WF, Lewis SL, Lloyd J, Monteagudo A, Neill DA, Patiño S, Pitman NCA, Silva JNM, Martínez RV (2004) Variation in wood density determines spatial patterns in Amazonian forest biomass. *Glob Chang Biol* 10(5):545–562
- Bastin J, Berrahmouni N, Grainger A, Maniatis D, Mollicone D, Moore R, Patriarca C, Picard N, Sparrow B, Abraham EM, Aloui K, Atesoglu A, Attore F, Bassüllü Ç, Bey A, Garzuglia M, García-Montero LG, Groot N, Guerin G, Laestadius L, Lowe AJ, Mamane B, Marchi G, Patterson P, Rezende M, Ricci S, Salcedo I, Sanchez-Paus Diaz A, Stolle F, Surappaeva V, Castro R (2017) The extent of forest in dryland biomass. *Science* 356(6338):635–638
- Behera SK, Sahu N, Mishra AK, Bargali SS, Behera MD, Tuli R (2017) Aboveground biomass and carbon stock assessment in Indian tropical deciduous forest and relationship with stand structural attributes. *Ecol Eng* 99:513–524
- Bonino E (2006) Changes in carbon pools associated with a land use gradient in the Dry Chaco, Argentina. *For Ecol Manag* 223(1–3):183–189
- Borah M, Das D, Kalita J (2015) Tree species composition, biomass and carbon stocks in two tropical forest of Assam. *Biomass Bioenergy* 78:25–35
- Brown S, Lugo AE (1992) Aboveground biomass estimates for tropical moist forests of the Brazilian Amazon. *Interciencia* 17:8–18
- Brown S, Sathaye J, Cannell M, Kauppi PE (1996) Mitigation of carbon emission to the atmosphere by forest management. *Commonw For Rev* 75(1):80–91
- Brown S, Schroeder P, Birdsey R (1997) Aboveground biomass distribution of US eastern hardwood forests and the use of large trees as an indicator of forest development. *For Ecol Manag* 96(1–2):37–47
- Cairns MA, Brown S, Helmer EH, Baumgardner GA (1997) Root biomass allocation in the world's upland forests. *Oecologia* 111(1):1–11
- Canadell JG, Le Quéré C, Raupach MR, Field CB, Buitenhuis ET, Ciais P, Conway TJ, Gillett NP, Houghton RA, Marland G (2007) Contribution to accelerating atmospheric CO<sub>2</sub> growth from economic activity, carbon intensity and efficiency of natural sinks. *Proc Natl Acad Sci* 104(47):18866–18870
- Cao J, Zhang X, Deo R, Gong Y, Feng Q (2018) Influence of stand type and stand age on soil carbon storage in China's arid and semi-arid regions. *Land Use Policy* 78:258–265
- Champion HG, Seth SK (1968) A revised survey of the forest types of India. Govt. of India Publications, New Delhi, p 404
- Chaturvedi OP, Singh JS (1987) A quantitative study of the forest floor biomass, litter fall and nutrient return in a *Pinus roxburghii* forest in Kumaun Himalaya. *Vegetatio* 71(2):97–106
- Chaturvedi RK, Raghubanshi AS, Singh JS (2011) Carbon density and accumulation in woody species of tropical dry forest in India. *For Ecol Manag* 262(8):1576–1588
- Chave J, Coomes DA, Jansen S, Lewis SL, Swenson NG, Zanne AE (2009) Towards a worldwide wood economics spectrum. *Ecol Lett* 12(4):351–366
- Chen L, Liang M, Wang S (2016) Carbon stock density in planted versus natural *Pinus massoniana* forests in sub-tropical China. *Ann Forest Sci* 73(2):461–472
- Chhabra A, Palria S, Dadhwal VK (2002) Growing stock-based forest biomass estimate for India. *Biomass Bioenergy* 22(3):187–194
- Chhabra A, Palria S, Dadhwal VK (2003) Soil organic carbon pool in Indian forests. *For Ecol Manag* 173(1):187–199
- Chibbar RK (1985) Soils of Delhi and their management. In: Biswas BC, Yadav DS, Maheshwari S (eds) Soils of India and their management. Fertiliser Association of India, New Delhi, pp 72–86
- Clark DB, Clark DA (1996) Abundance, growth, and mortality of very large trees in neotropical lowland rain forest. *For Ecol Manag* 80(1–3):235–244
- Dai Z, Birdsey RA, Johnson KD, Dupuy JM, Hernandez-Stefanoni JL, Richardson K (2014) Modeling carbon stocks in a secondary tropical dry forest in the Yucatan Peninsula, Mexico. *Water Air Soil Pollut* 225:1925
- Dar JA, Sundarapandian S (2015) Variation of biomass and carbon pools with forest type in temperate forests of Kashmir Himalaya, India. *Environ Monit Assess* 187(2):55
- Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (2006) IPCC guidelines for national greenhouse gas inventories, volume - IV agriculture, forestry and other land use. Institute of Global Environmental Strategies (IGES), Hayama
- FAO (2005) Global forest resources assessment: progress towards sustainable forest management. In: Davis R (ed) Food and agriculture organization of the united extent of forest resources nations, Rome
- FSI (1996) Volume equations for forests of India, Nepal and Bhutan. Forest Survey of India, Ministry of Environment and Forests, Government of India, India
- FSI (2017) State of forest report. Forest Survey of India, Ministry of Environment and Forests, Government of India, India
- Gandhi DS, Sundarapandian S (2017) Soil carbon stock assessment in the tropical dry deciduous forest of the Sathanur reserve forest of Eastern Ghats, India. *J Sustainable For* 36(2):1–17
- Gaxiola A, Armesto JJ (2015) Understanding litter decomposition in semiarid ecosystems: linking leaf traits, UV exposure and rainfall variability. *Front Plant Sci* 6:40
- Gibbs HK, Brown S, Niles JO, Foley JA (2007) Monitoring and estimating tropical forest carbon stocks: making REDD a reality. *Environ Res Lett* 2(4):1–13
- Gray JM, Bishop TFA, Wilson BR (2015) Factors controlling soil organic carbon stocks with depth in eastern Australia. *Soil Sci Soc Am J* 79(6):1741–1751
- Hou L, Xi WM, Zhang SX (2015) Effect of understorey on a natural secondary forest ecosystem carbon budget. *Russ J Ecol* 46(1):51–58
- Hu Y, Su Z, Li W, Li J, Ke X (2015) Influence of tree species composition and community structure on carbon density in a subtropical forest. *PLoS One* 10(8):e0136984
- Huang J, Guan X, Ji F (2012) Enhanced cold-season warming in semi-arid regions. *Atmos Chem Phys* 12(12):5391–5398
- Huang J, Yu H, Guan X, Wang G, Guo R (2016) Accelerated dryland expansion under climate change. *Nat Clim Chang* 6:166–171
- IPCC (2006) IPCC (Intergovernmental Panel on Climate Change) guidelines for national greenhouse gas inventories. In: Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (eds) National greenhouse gas inventories programme, Institute For Global Environmental Strategies, Japan
- Johnson MG, Kern JS (2002) Quantifying the organic carbon held in forested soils of the United States and Puerto Rico. In: Kimble JM, Heath LS, Birdsey RA, Lal R (eds) The potential of U.S. forest soils to sequester carbon and mitigate the greenhouse effects. Lewis Publishers, Boca Raton
- Joshi RK, Dhyani S (2018) Biomass, carbon density and diversity of tree species in tropical dry deciduous forests in Central India. *Acta Ecol Sin* <https://doi.org/10.1016/j.chnaes.2018.09.009>
- Júnior LRP, Andrade EMD, Palácio HADQ, Raymer PCL, Ribeiro FJC, Pereira FJS (2016) Carbon stocks in a tropical dry forest in Brazil. *Rev Ciênc Agron* 47(1):32–40
- Kauppi PA, Birdsey RA, Pan Y, Ihalainen A, Nöjd P, Lehtonen A (2015) Effect of land management on large trees and carbon stock. *Biogeosciences* 12:855–862
- Knight DH (1975) A phytosociological analysis of species-rich tropical forest on Barro Colorado Island, Panama. *Ecol Monogr* 45(3):259–284
- Köhl M, Neupane PR, Lotfiomran N (2017) The impact of tree age on biomass growth and carbon accumulation capacity: a retrospective analysis using tree ring data of three tropical tree species grown in natural forests of Suriname. *PLoS One* 12(8):e0181187
- Lal M, Singh R (2000) Carbon sequestration potential of Indian forests. *Environ Monit Assess* 60(3):315–327
- Lal R (2005) World crop residues production and implication of its use as a biofuel. *Environ Int* 31(4):575–586
- Malagnoux M, Sène EH, Atzmon N (2007) Forests, trees and water in arid lands: a delicate balance. *Unasyla* 58:24–29
- Manhas RK, Negi JDS, Kumar R, Chauhan PS (2006) Temporal assessment of growing stock, biomass and carbon stock of Indian forests. *Clim Chang* 74(1–3):191–221
- Meena A, Hanief M, Bidalia A, Rao KS (2016) Structure, composition and diversity of tree strata of semi-arid forest community in Delhi, India. *Phytomorphology* 66(3–4):95–102
- Meza FJ, Montes C, Bravo-Martínez F, Serrano-Ortiz P, Kowalski AS (2018) Soil water content effects on net ecosystem CO<sub>2</sub> exchange and actual

- evapotranspiration in a Mediterranean semiarid savanna of Central Chile. *Sci Rep* 8:8570
- Mills AJ, O'Connor TG, Donaldson JS, Fey MV, Skowno AL, Sigwela AM, Lechmere-Oertel RG, Bosenberg JD (2005) Ecosystem carbon storage under different land uses in three semi-arid shrublands and a mesic grassland in South Africa. *S Afr J Sci* 22(3):183–190
- Misra R (1968) Ecology work book. Calcutta, Oxford and IBH Publishing company, New Delhi, p 242
- Mohanraj R, Saravanan J, Dhanakumar S (2011) Carbon stock in Kolli forests, Eastern Ghats (India) with emphasis on aboveground biomass, litter, woody debris and soils. *IFOREST* 4:61–65
- Muhati GL, Olago D, Olaka L (2018) Quantification of carbon stocks in Mount Marsabit Forest Reserve, a sub-humid montane forest in northern Kenya under anthropogenic disturbance. *Glob Ecol Conserv* 14:e00383
- Naveenkumar J, Arunkumar KS, Sundarapandian SM (2017) Biomass and carbon stocks of a tropical dry forest of the Javadi Hills, Eastern Ghats, India. *Carbon Manag* 8(5–6):351–361
- Ovington JD (1956) The composition of tree leaves. *Forestry* 29(1):22–28
- Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, Phillips OL, Shvidenko A, Lewis SL, Canadell JG, Ciais P, Jackson RB, Pacala SW, McGuire AD, Piao S, Rautiainen A, Sitch S, Hayes D (2011) A large and persistent carbon sink in the world's forests. *Science* 333(6045):988–993
- Pande PK, Bisht APS, Sharma SC (1988) Comparative vegetation analysis of some plantation ecosystems. *Ind For* 114(7):379–388
- Post WM, Kwon KC (2000) Soil carbon sequestration and land use change: processes and potential. *Glob Chang Biol* 6(3):317–327
- Raghubanshi AS (1991) Dynamics of soil biomass C, N, and P in a dry tropical forest in India. *Biol Fertil Soils* 12(1):55–59
- Rajput SS, Shukla NK, Gupta VK, Jain JD (1996) Timber mechanics: strength classification and grading of timber. Indian Council of Forestry Research and Education, New Forest, Dehradun, p 189
- Ramachandran A, Jayakumar S, Haroon RM, Bhaskaran A, Arockiasamy DI (2007) Carbon sequestration: estimation of carbon stock in natural forests using geospatial technology in the Eastern Ghats of Tamil Nadu, India. *Curr Sci* 92(3):323–331
- Ravindranath NH, Ostwald M (2008) Methods for estimating above ground biomass. In: Ravindranath NH, Ostwald M (eds) Carbon inventory methods: handbook for greenhouse gas inventory, carbon mitigation and round wood production projects, Springer Science + Business Media B.V, Netherlands
- Ravindranath NH, Somasekhar BS, Gadgil M (1997) Carbon flows in Indian forest. *Clim Chang* 35(3):297–320
- Sahu SC, Suresh HS, Ravindranath NH (2016) Forest structure, composition and above ground biomass of tree community in tropical dry forests of Eastern Ghats, India. *Not Sci Biol* 8(1):125–133
- Salunkhe O, Khare PK, Kumari R, Kumari R, Khan ML (2018) A systematic review on the aboveground biomass and carbon stocks of Indian forest ecosystems. *Ecol Process* 7:17
- Santos LTD, Marra DM, Trumbore S, Camargo PB, Chambers JQ, Negrón-Juárez RI, Lima AJN, Ribeiro GHPM, Santos JD, Higuchi N (2016) Windthrows increase soil carbon stocks in a Central Amazon forest. *Biogeosciences* 12:19351–19372
- Sharma CM, Baduni NP, Gairola S, Ghildiyal SK, Suyal S (2010) Tree diversity and carbon stocks of some major forest types of Garhwal Himalaya, India. *For Ecol Manag* 260(12):2170–2179
- Singh J (1990) Nature and distribution of soil humic substances of some plant communities at Ooty, Nilgiri. *Myoforestry* 26:143–147
- Singh K, Chand P (2012) Above-ground tree outside forest (TOF) phytomass and carbon estimation in the semi-arid region of southern Haryana: a synthetic approach of remote sensing and field data. *J Earth Syst Sci* 121(6):1469–1482
- Sinha GN (2014) An introduction to the Delhi ridge. Department of Forests and Wildlife, Govt. of NCT of Delhi, New Delhi, India, p Xxiv + 154
- Solomon N, Birhane E, Tadesse T, Treydte AC, Meles K (2017) Carbon stocks and sequestration potential of dry forests under community management in Tigray, Ethiopia. *Ecol Process* 6:20
- Solomon N, Pabi O, Annang T, Asante IK, Birhane E (2018) The effects of land cover change on carbon stock dynamics in a dry Afromontane forest in northern Ethiopia. *Carbon Balance Mange* 13:14
- Stephenson NL, Das AJ, Condit R, Russo SE, Baker PJ, Beckman NG, Coomes DA, Lines ER, Morris WK, Rüger N, Álvarez E, Blundo C, Bunyavejchewin S, Chuyong G, Davies SJ, Duque Á, Ewango CN, Flores O, Franklin JF, Grau HR, Hao Z, Harmon ME, Hubbell SP, Kenfack D, Lin Y, Makana JR, Malizia A, Malizia LR, Pabst RJ, Pongpattananurak N, Su SH, Sun IF, Tan S, Thomas D, van Mantgem PJ, Wang X, Wiser SK, Zavalá MA (2014) Rate of tree carbon accumulation increases continuously with tree size. *Nature* 507:90–93
- Taylor AR, Wang JR, Chen HY (2007) Carbon storage in a chronosequence of red spruce (*Picea rubens*) forests in central Nova Scotia, Canada. *Can J For Res* 37(11):2260–2269
- Vayreda J, Gracia M, Canadell JG, Retana J (2012) Spatial patterns and predictors of forest carbon stocks in western Mediterranean. *Ecosystems* 15(8):1258–1270
- Wagner B, Liang E, Li X, Dulamsuren C, Leuschner C, Hauck M (2015) Carbon pools of semi-arid *Picea crassifolia* forests in the Qilian Mountains (north-eastern Tibetan Plateau). *For Ecol Manag* 343:136–143
- Wei Y, Li M, Chen H, Lewis BJ, Yu D, Zhou W, Fang X, Zhao W, Dai L (2013) Variation in carbon storage and its distribution by stand age and forest type in boreal and temperate forests in Northeast China. *PLoS One* 8(8):e72201
- Westlake DF (1963) Comparison of plant productivity. *Biol Rev* 38(3):385–425
- Yanai RD, Stehman SV, Arthur MA, Prescott CE, Friedland AJ, Siccama TG, Binkley D (2003) Detecting change in forest floor carbon. *Soil Sci Soc Am J* 67(5):1583–1593
- Yang YS, Guo JF, Chen G, Xie J, Gao R, Zhen L, Zhao J (2005) Carbon and nitrogen pools in Chinese fir and evergreen broadleaved forests and changes associated with felling and burning in mid-subtropical China. *For Ecol Manag* 216(1–3):216–226
- Yue J, Guan J, Deng L, Zhang J, Li G, Du S (2018) Allocation pattern and accumulation potential of carbon stock in natural spruce forests in Northwest China. *PeerJ* 6:e4859
- Zanne AE, Lopez Gonzalez G, Comes DA, Ilic J, Janson S, Lewis SL, Miller RB, Swenson NG, Wiemann MC, Chave J (2009) Global wood density database. Dryad digital repository
- Zhang D, Hui D, Luo Y, Zhou G (2008) Rates of litter decomposition in terrestrial ecosystems: global patterns and controlling factors. *J Plant Ecol* 1(2):85–93
- Zhang Q, Wang C (2010) Carbon density and distribution of six Chinese temperate forests. *Sci Chin* 53(7):831–840
- Zhang Y, Gub F, Liu S, Liu Y, Li C (2013) Variations of carbon stock with forest types in subalpine region of southwestern China. *For Ecol Manag* 300:88–95

**Submit your manuscript to a SpringerOpen® journal and benefit from:**

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)

---