

RESEARCH

Open Access



Using hierarchical spatial models to assess the occurrence of an island endemism: the case of *Salamandra corsica*

Daniel Escoriza^{1*}  and Axel Hernandez²

Abstract

Background: Island species are vulnerable to rapid extinction, so it is important to develop accurate methods to determine their occurrence and habitat preferences. In this study, we assessed two methods for modeling the occurrence of the Corsican endemic *Salamandra corsica*, based on macro-ecological and fine habitat descriptors. We expected that models based on habitat descriptors would better estimate *S. corsica* occurrence, because its distribution could be influenced by micro-environmental gradients. The occurrence of *S. corsica* was modeled according to two ensembles of variables using random forests.

Results: *Salamandra corsica* was mainly found in forested habitats, with a complex vertical structure. These habitats are associated with more stable environmental conditions. The model based on fine habitat descriptors was better able to predict occurrence, and gave no false negatives. The model based on macro-ecological variables underestimated the occurrence of the species on its ecological boundary, which is important as such locations may facilitate interpopulation connectivity.

Conclusions: Implementing fine spatial resolution models requires greater investment of resources, but this is advisable for study of microendemic species, where it is important to reduce type II error (false negatives).

Keywords: Amphibian, Corsica, Mediterranean islands, Microhabitat, Salamander

Introduction

Macro-ecological models are popular for evaluating ecological and evolutionary hypotheses for vertebrates. This approach is commonly used because the free access to databases, including Worldclim and the Global Biodiversity Information Facility, enables the evaluation of hypotheses involving large taxonomic groups over broad geographical areas (Buckley and Jetz 2007; Araújo et al. 2008; Gregory et al. 2009). Studies involving the use of fine spatial scale descriptors are expensive in terms of resources required per species and site, which typically restricts their application to a limited number of species in small regions (Freemark and Merriam 1986; Williams et al. 2002). However, in some cases (e.g., forest ectothermic vertebrates) macroclimatic models cannot adequately describe a species niche, leading to erroneous extrapolations (Scheffers et al. 2014a). For this

reason, studies evaluating the niche of ectothermic vertebrates associated with forests should include parameters that describe the habitats in finer detail (Petranka et al. 1993; Rödel and Ernst 2004; Escoriza et al. 2018).

Urodeles are particularly sensitive to environmental conditions because they have narrow ranges of thermal/humidity tolerance (Hutchison 1961; Spotila 1972). Many species of terrestrial salamander are habitat specialists, confined to thermally buffered microhabitats including riparian forests, caves, or fissured substrates (Salvidio 1998; Johnston and Frid 2002; Godmann et al. 2016). This suggests that terrestrial salamanders are useful for testing the effectiveness of ecological models constructed using sets of variables covering different spatial scales. In addition, a large proportion of salamander species are threatened globally, by habitat loss and the expansion of pathogens (Rovito et al. 2009; Martel et al. 2013). For this reason to have accurate tools for describing their habitat preferences could aid their conservation, including defining optimal survival habitats so that

* Correspondence: daniel_escoriza@hotmail.com

¹GRECO, Institute of Aquatic Ecology, University of Girona, 17071 Girona, Spain

Full list of author information is available at the end of the article

such areas can be included in natural reserves (Esselman and Allan 2011). Terrestrial salamanders are poor terrestrial dispersers and frequently the territory of adults is close to breeding habitats (Bar-David et al. 2007). It can be assumed that the optimal habitats for the survival of populations are those that include suitable habitats for reproduction and for surface activity of adults.

In this study, we evaluated the performance of two modeling methods for predicting the occurrence of an island salamander, *Salamandra corsica*. This species is endemic to Corsica but distributed discontinuously, and the factors explaining the pattern of occurrence are not fully understood (Lanza et al. 2007; Bosc and Destandau 2012). The fact that this species is confined to a geographically small region provides the opportunity to characterize most of its distributional range at fine spatial resolution, in contrast to other species of the Mediterranean salamander. In addition, it is a priority to understand the ecological requirements of *S. corsica*, because island species are particularly prone to rapid extinction because of the accidental translocation of pathogens or alien species (Sax and Gaines 2008).

Corsica has a very heterogeneous landscape, with major topographic gradients in climate and plant communities (Jeanmonod and Gamisans 2007). Characterizing these communities can be useful in describing the local landscape complexity, because plants respond at fine scale and in competitive ways to subtle changes in temperature, moisture, substrate chemistry, and anthropogenic disturbance (Beatty 1984; Wilson and Keddy 1986). Moreover, plants constitute shelter and food for animals, so their importance is not limited to reflecting the abiotic background (Garden et al. 2007). For these reasons, the gradients of vegetation have a marked influence on the organization of zoological assemblages (Bersier and Meyer 1994).

In this study, using macro- and micro-ecological data, we investigated the capacity of two statistical models to determine and predict the occurrence of the endemic *S. corsica* found in Corsica Island, western Mediterranean. The macro-ecological variables described the landscape and climate at coarse spatial resolution, and were obtained from freely available databases (Chander et al. 2009; Fick and Hijmans 2017). The micro-ecological variables described the fine composition and structure of the terrestrial habitats involved, particularly focusing on those parameters that determine, or are associated with, thermal/humidity buffers (Johnston and Frid 2002); these were characterized in situ in each locality where the target species occurred. We predicted that those models that included habitat descriptors would better estimate *S. corsica* occurrence, because the distribution of the species could be influenced by micro-gradients that are not described

by large-scale spatial models (Scheffers et al. 2014b). We expected this pattern because this salamander appears discontinuously in the meso-Mediterranean climate belt, suggesting the use of habitats associated with very specific topographic and vegetation profiles (Lanza et al. 2007).

Methods

Study area and surveys

The study region was the island of Corsica (8680 km²), which is located in the western Mediterranean. Two thirds of the surface of the island is mountainous, with a maximum elevation of 2706 m at Monte Cinto (van Geldern et al. 2014). The climate of the island is *Csa* type (Köppen system) on the peri-coastal plains, but the *Csb* type predominates inland, from elevations of 400 m (Peel et al. 2007). Most of the island receives moderate to high rainfall, from 774 mm year⁻¹ at Vizzavona (974 m) to 547 mm year⁻¹ at Bonifacio (63 m). The natural vegetation is dominated by meso-Mediterranean forests (*Quercus ilex/Q. suber*) and evergreen shrubs ('maquis'; 0–700 m), supra-Mediterranean forests (*Pinus nigra/Quercus pubescens*; 700–1300 m) and oro-Mediterranean and temperate forests (*Fagus sylvatica/P. nigra*; 1000–1800 m; IFN (Inventaire Forestier National) 2006; Gamisans 2010). Above 1600–1800 m, sub-alpine fir forests (*Abies alba*) and shrubs predominate (Gamisans 2010).

Salamandra corsica is one of two species of urodeles that occur on Corsica (Delaugerre and Cheylan 1992). This salamander is endemic to the island, where it occurs from 50 to 1890 m, although it is rare below 400–500 m, where it is mostly confined to humid forests (Lanza et al. 2007; Lescure and de Massary 2012). According to the IUCN, this species is not threatened but its ecological requirements are poorly known, and it could be adversely affected by habitat disturbance and emerging diseases (Miaud et al. 2009; Bosc and Destandau 2012). We surveyed *S. corsica* over a year from September 2014 to September 2015, and for six shorter periods of 10 days in October, November, February, March, and May in the years 2006 and 2018. Surveying in different months and years reduced the uncertainty associated with interannual variability in the use of aquatic habitats by amphibians under unstable hydrological regimes (Gómez-Rodríguez et al. 2010). Area-constrained surveys (30 × 30 m) were conducted by assessing ponds, springs, and streams for larvae (Duguet and Melki 2003), and assessing terrestrial habitats on foot for adults during the day and at night in rainy weather (Hernandez et al. 2018). We lifted rock and logs under which larvae and adult salamanders usually seek refuge.

The surveyed sites were classified according to the presence or absence of *S. corsica*. Assessing for the

occurrence of a terrestrial salamander at a given site can be difficult because they are elusive or confined to microhabitat patches (Warburg 1986; Surasinghe and Baldwin 2015). Sites for which we determined that the species was absent were defined as those outside the convex hull generated by the known species distribution range (Delaugerre and Cheylan 1992; Bosc and Destandau 2012), and where *S. corsica* was not detected in at least two surveys conducted in different years and seasons (Fig. 1). The combination of several survey methods (i.e., terrestrial and aquatic habitats) under different environmental conditions at different periods of the year reduces the probability of including false absences in the survey data (Williams and Berkson 2004).

Data collection

Both presence and absence sites were characterized using environmental variables that described climate and landscape features important in the natural history of terrestrial salamanders (Kozak and Wiens 2010; Escoriza and Ben Hassine 2017). The climate was characterized by the maximum temperature in July and the minimum temperature in January ($^{\circ}\text{C}$), water vapor pressure in July

(kPa), and mean annual precipitation (mm year^{-1}) (Hijmans et al. 2005; Fick and Hijmans 2017). Land cover was characterized using a vegetation index (enhanced vegetation index, EVI; Huete et al. 2002). The EVI data were obtained from Landsat 8 (32-day composite) for the month of July (Chander et al. 2009). This vegetation index is less sensitive to random environmental noise than the normalized difference vegetation index (NDVI), and describes variations in the biomass and in the architecture of the forest canopy (Pettorelli et al. 2005). The topography descriptors included altitude and the terrain ruggedness index, an estimator of topographic heterogeneity (Riley et al. 1999). The terrain ruggedness index was obtained from a digital elevation model having a resolution of 250 m (Jarvis et al. 2008). Elevation was obtained in situ using a global positioning system at an accuracy of ± 15 m (Garmin Etrex 10; Garmin Ltd., Olathe, KS, USA). GIS environmental variables data were extracted using QGIS v. 3.4.1 (QGIS Development Team 2018).

We also characterized the composition and structure of plant communities at the study sites (James and Wamer 1982). The richness of plant species in Corsica is too high (2858 species; Jeanmonod and Gamisans 2007)




Fig. 1 Map of the study region. The shaded polygon shows the distribution of *Salamandra corsica* according to the IUCN (International Union for Conservation of Nature and Natural Resources) (2018). Blue triangles, presence sites; orange circles, absence/undetected presence sites

to be useful in a statistical model. For this reason, we summarized plant diversity by grouping species into functional taxonomic types, widely used in standard botanical classifications (Thorogood 2018). These groups were (i) height type: trees (species having a maximum height > 5 m), shrubs (1–5 m), sub-shrubs (< 1 m), lianas (climbing plants), and perennial grasses (plants lacking woody stems and having a lifespan > 2 years); (ii) leaf type: aciculifolious (conifers), deciduous, and perennial; and (iii) taxonomic groups: Spermatophyta (including all previous subgroups), Lycopodiophyta (lycophytes), and Pteridophyta (equisetums and ferns). In addition, we included a group comprising naturalized plants (alien plants of spontaneous growth, following Gamisans 2010, 2014) as bio-indicators of anthropic disturbance (Larson et al. 2001). Data regarding plant composition were collected by sampling along three transects, each of 30 m in length radiating in north, southeast, and southwest directions (respectively) from a central point of occurrence of *S. corsica* (Guénette and Villard 2005). The relative importance of these groups to the vertical structure of the plant community was estimated by multiplying the number of stems of the species by their average heights. Tree density combined size estimations

are commonly used in forest inventories to assess the dominance of tree species (McIntyre et al. 2015). The height was estimated by measuring average-sized specimens, using images calibrated in ImageJ 1.52 (Rasband 2018). The dataset supporting the conclusions of this article is included within the article and its Additional file 1.

Data analyses

Correlations involving all variables were tested. Those that showed a correlation value > 0.75 were removed from subsequent analyses (Griffith et al. 2003). The remaining variables were normalized. The relative position of the presence and absence sites within the environmental gradient was visualized using principal component analysis (PCA). We assessed whether the two groups of sites differed significantly in their environmental conditions, applying a PERMANOVA test with the matrix based on Euclidean distances (one fixed factor = presence/absence) (Anderson 2001). These analyses were carried out using the package PRIMER-E (Primer-E Ltd., Plymouth) and the software 'pca3d' (Weiner 2017) for R v. 3.5.3 (R Development Core Team 2018).

The significant associations and the sense (positive or negative) of the associations between the presence of *S.*

Table 1 Descriptive statistics (mean \pm standard error) of the environmental variables. n = number of sites

Variables	Scale	Presence	Absence
n		35	31
Annual precipitation (mm year ⁻¹)	1000 m	728 \pm 11	617 \pm 13
Enhanced Vegetation Index July	500 m	0.57 \pm 0.03	0.40 \pm 0.02
Maximum temperature July (°C)	1000 m	24.6 \pm 0.4	27.8 \pm 0.2
Minimum temperature January (°C)	1000 m	1.1 \pm 0.3	5.0 \pm 0.2
Water vapor pressure July (kPa)	1000 m	1.7 \pm 0.0	1.9 \pm 0.0
Elevation (masl)	15 m	590 \pm 59	54 \pm 10
Terrain ruggedness	250 m	161 \pm 11	47 \pm 6
Aciculifolious bush	30 m	10 \pm 7	28 \pm 17
Aciculifolious tree	30 m	165 \pm 56	19 \pm 10
Alien plants %	30 m	9.5 \pm 3.1	3.8 \pm 2.0
Deciduous broad-leaved bush	30 m	11 \pm 2	11 \pm 3
Deciduous broad-leaved tree	30 m	416 \pm 55	38 \pm 13
Deciduous liana	30 m	21 \pm 7	0.3 \pm 0.2
Equisetum	30 m	0.1 \pm 0.1	0.1 \pm 0.1
Fern	30 m	27 \pm 5	2 \pm 1
Lycophyte	30 m	0.2 \pm 0.2	0.0
Perennial broad-leaved bush	30 m	154 \pm 22	240 \pm 33
Perennial broad-leaved tree	30 m	287 \pm 36	242 \pm 38
Perennial grass	30 m	42 \pm 7	42 \pm 12
Perennial liana	30 m	72 \pm 10	56 \pm 12
Sub-shrub	30 m	31 \pm 6	49 \pm 6




Table 2 Binomial generalized linear models results assessing the environmental variation between presence/absence sites. The model-averaged importance for the variables present in the 100 best models is shown. % models, percentage of candidate models where the variable was included

Variable	Univariate estimate	Univariate <i>P</i>	% models	Importance
Elevation	12.104	0.002	71	0.738
Deciduous broad-leaved tree	3.698	0.0004	51	0.483
Fern	4.137	0.002	43	0.461
Deciduous liana	18.896	0.003	28	0.233
Maximum temperature July	−6.308	0.0004	22	0.210
Enhanced Vegetation Index July	1.156	0.0009	14	0.180
Perennial broadleaved tree	0.223	0.379	18	0.162
Aciculifolious bush	−0.277	0.341	18	0.149
Deciduous broad-leaved bush	0.019	0.940	13	0.142
Aciculifolious tree	1.528	0.076	13	0.100
Perennial liana	0.258	0.322	13	0.094
Perennial grass	0.003	0.989	11	0.093
Perennial broad-leaved bush	−0.571	0.037	11	0.093
Terrain ruggedness	3.337	0.00003	10	0.073
Sub-shrub	−0.572	0.050	9	0.064
Annual precipitation	1.944	0.00001	7	0.057

corsica and the environmental descriptors were estimated separately using univariate binomial generalized linear models (GLMs). A univariate method was used to test the sense and the statistical significance (alpha level = 0.05) of the associations between the dependent variable and predictors, not biased by variable correlations (Dormann et al. 2013). The relative importance of the pooled variables was evaluated using automated model selection and multi-model inference procedures for GLMs (Calcagno and de Mazancourt 2010). The best candidate models were obtained using a holistic selection, and the models were ranked using the small-sample corrected Akaike information criterion (AICc; Burnham and Anderson 2002). We assessed the statistical importance of each variable according to its model-averaged weighting in the 100 best models. These analyses were conducted using the package ‘glmulti’ (Calcagno 2015) in R.

The efficiency of the model in determining the presence of *S. corsica* at the study sites was assessed using random forests (Cutler et al. 2007). This classification method has been shown to be superior to other classification algorithms, and is less prone to overfit when using high dimensional datasets (Hastie et al. 2009; Broennimann et al. 2012). We built two models, one including only macro-ecological parameters (hereafter $\hat{y}1$ model) and having a spatial resolution of 250–1000 m (except for elevation, which had a spatial resolution of 15 m). The other model was built by combining macro-ecological and microhabitat parameters (hereafter $\hat{y}2$ model).

We set the random forest parameters for classification, and used 10,000 training trees (Liaw and Wiener 2002). The optimum number of variables randomly sampled at each split was estimated by the decrease in the out-of-bag (OOB) error, using the ‘tuneRF’ function implemented in the ‘randomForest’ package (Liaw and Wiener 2002). We assessed model performance based on several estimators that evaluated the efficiency, rate, and type of errors in the discriminatory capacity of binary models, including the OOB error, the area under the receiver-operating characteristic curve (AUC), and the rates of false positives (FP) and false negatives (FN) (Lalkhen and McCluskey 2008; Mwitondi 2013). The OOB is the misclassification rate of the random forest model estimated for the training data, with higher values indicating models having lower classification accuracy (Mwitondi 2013). The AUC facilitated evaluation of the predictive efficiency of the model; its values ranged from 0.5 for models having predictive capacity similar to chance, to 1.0 for models having perfect predictive ability (Araújo et al. 2005). We evaluated variable importance of each variable based on the mean decrease accuracy (MDA; loss of model accuracy if the variable is excluded from the model; Guo et al. 2016). These analyses were carried out with the package ‘randomForest’ (Liaw and Wiener 2002) and ‘ROCR’ (Sing et al. 2015) in R.

Results

Salamandra corsica was found at 35 sites, encompassing most of the island of Corsica (Fig. 1). The species

Table 3 Predicted probabilities for the presence/absence sites using random forest classification. Observed (+) = *Salamandra corsica* presence site; Observed (−) = *Salamandra corsica* absence site. $\hat{Y}1$ = macro-ecological model; $\hat{Y}2$ = mixed model. Datum for all coordinates, WGS84. Elevation, masl

Locality	Latitude	Longitude	Elevation	Vegetation	Observed	$\hat{Y}1$	$\hat{Y}2$
Source de Pizzolo	42.07	9.14	1722	Deciduous forest	+	0.999	0.946
Bocca di Sorba	42.14	9.19	1338	Coniferous forest	+	0.978	0.847
Palneca	41.99	9.18	1112	Deciduous forest	+	0.999	0.995
Barrage de l'Ospédale	41.66	9.20	955	Coniferous forest	+	0.988	0.925
Bocca d'Illarata	41.70	9.21	954	Coniferous forest	+	0.949	0.816
Punta di u Diamante	41.69	9.21	953	Coniferous forest	+	0.998	0.891
Bocca di Vizzavona	42.11	9.13	839	Deciduous forest	+	0.999	0.977
Bastelica	41.98	9.04	826	Deciduous forest	+	0.998	0.998
Quenza	41.77	9.13	768	Evergreen forest	+	0.634	0.732
Campodonico	42.38	9.35	711	Deciduous forest	+	1.000	0.974
Muracciole	42.17	9.18	708	Evergreen forest	+	1.000	0.903
Bavella	41.81	9.25	673	Mixed forest	+	0.998	0.969
Perelli	42.32	9.39	644	Deciduous forest	+	0.998	0.954
Venaco	42.23	9.17	636	Deciduous forest	+	1.000	0.961
Casalabriva	41.74	8.93	587	Evergreen forest	+	0.997	0.967
Rivière du Vecchio	42.19	9.16	575	Evergreen forest	+	1.000	0.939
Chidazzu	42.23	8.77	572	Evergreen forest	+	0.986	0.995
Palasca	42.58	9.04	557	Evergreen forest	+	0.989	0.953
Bisene	41.65	9.09	532	Evergreen forest	+	0.870	0.918
Bocca di Sorru	42.17	8.85	499	Evergreen forest	+	1.000	0.856
Levie	41.69	9.13	463	Evergreen forest	+	0.999	0.983
Novale	42.30	9.43	463	Evergreen forest	+	1.000	0.944
Col de Bacinu	41.61	9.16	447	Evergreen forest	+	0.990	0.911
Cauro	41.91	8.92	381	Evergreen forest	+	0.772	0.830
Chiatra	42.28	9.47	350	Evergreen forest	+	0.999	0.993
Chialza	41.65	9.05	344	Evergreen forest	+	0.864	0.912
Zoza	41.73	9.08	317	Evergreen forest	+	0.992	0.926
Lapedina	42.85	9.42	293	Evergreen forest	+	0.983	0.893
Oreta	42.84	9.41	276	Evergreen forest	+	0.959	0.923
Casta	42.66	9.19	256	Maquis	−	0.960	0.567
Moltifao	42.48	9.15	251	Evergreen forest	+	0.385	0.624
Bocca Albitrina	41.60	8.93	240	Evergreen forest	+	0.943	0.536
Girolata	42.34	8.66	215	Maquis	+	0.939	0.669
Porto	42.25	8.67	167	Evergreen forest	+	0.865	0.932
Appietto	42.00	8.73	158	Maquis	+	0.531	0.751
Chiova d'Asinu	41.48	9.24	158	Maquis	−	0.833	0.302
Patrimonio	42.70	9.37	139	Evergreen forest	+	0.437	0.654
Figari	41.47	9.12	99	Maquis	−	0.073	0.185
Oletta	42.61	9.29	99	Maquis	−	0.012	0.206
Cargèse	42.17	8.62	97	Maquis	−	0.152	0.287
Cuo	41.53	9.17	89	Evergreen forest	−	0.005	0.033
Alzone	41.86	8.80	88	Evergreen forest	−	0.017	0.043

Table 3 Predicted probabilities for the presence/absence sites using random forest classification. Observed (+) = *Salamandra corsica* presence site; Observed (–) = *Salamandra corsica* absence site. $\hat{Y}1$ = macro-ecological model; $\hat{Y}2$ = mixed model. Datum for all coordinates, WGS84. Elevation, masl (Continued)

Locality	Latitude	Longitude	Elevation	Vegetation	Observed	$\hat{Y}1$	$\hat{Y}2$
Serraggia	41.51	8.94	88	Evergreen forest	–	0.004	0.041
Tizzano	41.55	8.89	88	Maquis	–	0.000	0.006
Pont de Teghia	42.26	9.50	85	Cultivated field	–	0.004	0.038
Prigugio	42.51	8.78	77	Maquis	–	0.007	0.035
Cavallo Morto	41.40	9.17	69	Maquis	–	0.055	0.135
Fossi	41.65	9.32	62	Maquis	–	0.022	0.015
Bartaccia	41.67	8.91	54	Maquis	–	0.000	0.026
Acqua Longa	41.95	8.78	51	Suburban field	–	0.001	0.046
Molini	41.85	8.81	50	Maquis	–	0.017	0.033
Campo al Quarcio	42.14	9.45	34	Cultivated field	–	0.000	0.030
Aristone	42.06	9.44	21	Evergreen forest	–	0.000	0.230
Ghisonaccia	42.02	9.40	21	Cultivated field	–	0.000	0.080
Caramontinu	41.69	9.37	19	Evergreen forest	–	0.000	0.299
Santa Giulia	41.53	9.26	17	Maquis	–	0.000	0.006
La Tonnara	41.42	9.10	15	Maquis	–	0.104	0.157
Gurgazu	41.39	9.23	14	Maquis	–	0.085	0.127
Fromontica	42.67	9.28	7	Coastal habitat	–	0.038	0.135
Porto–Vecchio	41.58	9.30	7	Coastal habitat	–	0.000	0.012
Losari	42.62	9.00	6	Cultivated field	–	0.005	0.084
Etang de Biguglia	42.65	9.44	5	Wetland	–	0.005	0.057
Portigliolo	41.64	8.86	4	Coastal habitat	–	0.000	0.032
Sagone	42.11	8.69	4	Cultivated field	–	0.007	0.060
Favarella	41.72	8.84	4	Cultivated field	–	0.019	0.076
Aléria	42.11	9.55	1	Coastal habitat	–	0.064	0.096

typically occurs in elevated habitats having high rainfall and moderate temperatures (Table 1). In the sites where the species was present, the vegetation had a complex vertical structure: high canopies of broadleaf/aciculifolious trees (approximately 10–30 m height), with a mid-understory of deciduous bushes (approximately 2–5 m height) and a low understory of lianas, ferns, and perennial grasses (approximately 1 m height) (Table 1 and Additional file 1).

Correlation matrices indicated that the water vapor pressure in July and the minimum temperature in January were highly correlated with the maximum temperature in July ($r > 0.75$). Thus, the variables temperature and water vapor pressure were removed from subsequent analyses. The PCA showed that the presence and absence sites were distributed separately along a multidimensional environmental gradient (explained variance = 28%; Fig. 2). The first axis of the PCA (explained variance = 11.02%) was negatively correlated with the elevation and was positively correlated with temperatures (Fig. 2). The second axis of the PCA (explained variance = 9.24%) was positively

correlated with aciculifolious trees and was negatively correlated with perennial liana (Fig. 2). The third axis of the PCA (explained variance = 7.89%) was positively correlated with perennial grass and was negatively correlated with perennial broadleaved trees (Fig. 2). The PERMANOVA test confirmed that both groups of sites differed statistically (Pseudo- $F = 13.147$, $P = 0.0001$).

The GLM analyses showed that the topography (elevation: positive association with *S. corsica* occurrence), vegetation structure (proportion of deciduous broad-leaved trees, ferns, deciduous lianas: positive association with *S. corsica* occurrence), and temperature (negative association with *S. corsica* occurrence) were the major variables (present in at least the 22% of the best models) differentiating presence and absence sites (Table 2). All these associations were statistically significant (Table 2).

Both models generated using a random forest classifier correctly discriminated a large proportion of the study sites (Table 3). The $\hat{y}1$ model showed AUC = 0.984 and OOB = 6.06% and the $\hat{y}2$ model showed AUC = 0.999 and OOB = 1.52%. The $\hat{y}2$ model was superior to the $\hat{y}1$

model in predicting those sites having ecologically limiting conditions (i.e., within the meso-Mediterranean maquis and/or at low elevations; Table 3), with a lower false negative rate ($FN_{\hat{y}2} = 0.000$ versus $FN_{\hat{y}1} = 0.057$). Also, the $\hat{y}2$ model showed a lower false positive rate ($FP_{\hat{y}2} = 0.032$ versus $FP_{\hat{y}1} = 0.065$). The fine habitat descriptors also contributed substantially to discriminating between presence/absence localities in the $\hat{y}2$ model: deciduous broad-leaved tree (MDA = 48.12), fern (MDA = 36.80), and deciduous liana (MDA = 30.15) (Table 4).

Discussion

Our analyses showed that the model including micro-ecological and macro-ecological descriptors (the $\hat{y}2$ model) provided the best prediction of the occurrence of *S. corsica*. Although the difference in the performance of the models was relatively small, the $\hat{y}2$ model better predicted the occurrence of the species in environments at the edge of its distribution. This included low elevation meso-Mediterranean shrubland or ‘maquis,’ where the species is more localized in smaller populations. These populations could be important to the long-term survival of the species, acting as genetic corridors interconnecting mountain populations, as has been observed for other urodeles (Giordano et al. 2007). Furthermore, some peripheral amphibian populations show greater genetic

Table 4 Importance of the variables in the random forest model classification, showed as mean decrease accuracy (MDA; loss of model accuracy if the variable is excluded from the model, higher values indicate greater importance)

Variables	MDA
Elevation	86.16
Maximum temperature July	64.08
Terrain ruggedness	61.08
Deciduous broad-leaved tree	48.12
Annual precipitation	46.53
Fern	36.80
Deciduous liana	30.15
Enhanced Vegetation Index July	16.20
Aciculifolious tree	8.31
Perennial liana	8.02
Perennial broad-leaved bush	7.55
Sub-shrub	5.56
Alien plants %	4.30
Perennial grass	3.06
Deciduous broad-leaved bush	2.29
Perennial broad-leaved tree	0.42
Equisetum	0.00
Lycophyte	0.00
Aciculifolious bush	-2.65

diversity than those located in core areas, possibly because of the suboptimal environmental conditions (Munwes et al. 2010). However, no studies have evaluated whether this is also the case for *S. corsica*.

Consistent with the findings of Delaugerre and Cheylan (1992) and Lanza et al. (2007), our data showed that *S. corsica* occupies mainly temperate and supra-Mediterranean deciduous forests composed of beech (*F. sylvatica*) and chestnut (*Castanea sativa*), and meso-/oro-Mediterranean forests of black pine (*P. nigra*) and holm oak (*Q. ilex*), from 240 to 1722 masl. However, this salamander also occurred in meso-Mediterranean sclerophyllous shrubland, from 139 to 215 masl. The habitat of the sites where *S. corsica* was present differed structurally from those where it was absent. The two methods used to evaluate these differences showed that this could mainly be explained by the proportions of deciduous trees, ferns, and deciduous lianas. The deciduous broad-leaved trees included species with a height range of 15–30 m (mainly *Alnus glutinosa*, *C. sativa*, *F. sylvatica*, *Fraxinus ornus*, and *Q. pubescens*), which commonly formed a high and dense canopy poorly permeable to sunlight (Gálhidy et al. 2006). The group of deciduous lianas comprised species of the genera *Lonicera*, *Rosa*, and *Rubus*. All the species involved, together with ferns, are confined to habitats having high levels of edaphic moisture, such as Mediterranean riverine strips and/or humid montane forests (Gamisans 2010, 2014). Our data also showed that sites where *S. corsica* was present were also topographically heterogeneous. This heterogeneity, together with a dense forest canopy, is typically associated with more stable air temperatures and humidity (Suggitt et al. 2011; von Arx et al. 2012). This could explain why macroclimate models failed to predict the occurrence of *S. corsica* in the edge habitats, where it is mainly (or exclusively) located in or around closed forest ravines (Lanza et al. 2007).

Another aspect evaluated in this study was anthropogenic effects on the sites where *S. corsica* was present, determined by the presence of aliens in the plant community structure. The results indicated that *S. corsica* can occupy highly altered habitats in which alien species comprise up to 75.4% of the vegetation community. However, this was associated with the massive forests (about 24,600 ha; IFN (Inventaire Forestier National) 2006) of probably non-native chestnut on the island (Roces-Díaz et al. 2018). The chestnut forests typically comprise mature trees mixed with native trees and bushes, providing a vertical structure that mimics the natural deciduous forests (IFN (Inventaire Forestier National) 2006; Gamisans 2010), so the microclimatic conditions may also be similar. On the other hand, *S. corsica* was absent from other types of highly modified habitat, including

intensive agriculture and suburban habitats, where the vegetation structure does not resemble natural forest formations. Our findings, together with the data provided by the literature (Delaugerre and Cheylan 1992; Lanza et al. 2007), suggested that this species could avoid open human-disturbed landscapes.

Conclusions

Our findings showed that for describing the niche of a forest salamander, models including fine habitat descriptors are preferable to those based solely on macro-ecological parameters. This finding has important implications, particularly for the conservation of micro-endemic and threatened species, where it is crucial to minimize type II errors (sites are not protected, but include populations of the target species; Loiselle et al. 2003). Our method allows determining these sites with high probability of presence of the target species and therefore where to concentrate the survey effort to detect overlooked refuges or populations (Hughes et al. 2008).

The methodology used in this study is relatively simple to apply in the field, although it requires substantial survey effort (depending on the detectability of the species) and some botanical knowledge. The use of taxonomic and basic functional groups to describe plant associations enables this protocol to be used in any ecoregion worldwide, so its effectiveness could be tested in niche models generated for other species of forest-dwelling amphibian.

Additional file

Additional file 1: Data describing the characteristics of plant species found at the study sites. (XLSX 14 kb)

Acknowledgments

We thank the people of the Università di Corsica Pasquale Paoli (UCPP) for their assistance to conduct this research. We also thank Solomon Ayele Tadesse for his constructive comments on earlier drafts. This study was carried out respecting the legislation on wild fauna and flora of France. The authorizations were provided by the UCPP.

Funding

The authors have not received funds for this study.

Availability of data and materials

All data generated or analyzed during this study are included in this published article (and its supplementary information files).

Authors' contributions

AH and DE performed the field work. DE analyzed and interpreted the data. DE was a major contributor in writing the manuscript. AH substantially revised it. All authors read and approved the final manuscript

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹GRECO, Institute of Aquatic Ecology, University of Girona, 17071 Girona, Spain. ²Department of Environmental Sciences, University Pasquale Paoli, 20250 Corte, France.

Received: 1 March 2019 Accepted: 17 April 2019

Published online: 17 May 2019

References

- Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance. *Austral Ecol* 26:32–46. <https://doi.org/10.1111/j.1442-9993.2001.01070.pp.x>
- Araújo MB, Nogués-Bravo D, Diniz-Filho JAF, Haywood AM, Valdes PJ, Rahbek C (2008) Quaternary climate changes explain diversity among reptiles and amphibians. *Ecography* 31:8–15. <https://doi.org/10.1111/j.2007.0906-7590.05318.x>
- Araújo MB, Whittaker RJ, Ladle RJ, Erhard M (2005) Reducing uncertainty in projections of extinction risk from climate change. *Glob Ecol Biogeogr* 14: 529–538. <https://doi.org/10.1111/j.1466-822X.2005.00182.x>
- Bar-David S, Segev O, Peleg N, Hill N, Templeton AR, Schultz CB, Blaustein L (2007) Long-distance movements by fire salamanders (*Salamandra atra infraimaculata*) and implications for habitat fragmentation. *Isr J Ecol Evol* 53:143–159. <https://doi.org/10.1080/15659801.2007.10639579>
- Beatty SW (1984) Influence of microtopography and canopy species on spatial patterns of forest understory plants. *Ecology* 65:1406–1419. <https://doi.org/10.2307/1939121>
- Bersier LF, Meyer DR (1994) Bird assemblages in mosaic forests: the relative importance of vegetation structure and floristic composition along the successional gradient. *Acta Oecol* 15:561–576
- Bosc V, Destandau R (2012) Etude diagnostique de la Salamandre de Corse, *Salamandra corsica* (Savi, 1838). Office de l'Environnement de la Corse, Borgo
- Broennimann O, Fitzpatrick MC, Pearman PB, Petitpierre B, Pellissier L, Yoccoz NG, Thuiller W, Fortin MJ, Randin C, Zimmermann NE, Graham CH (2012) Measuring ecological niche overlap from occurrence and spatial environmental data. *Glob Ecol Biogeogr* 21:481–497. <https://doi.org/10.1111/j.1466-8238.2011.00698.x>
- Buckley LB, Jetz W (2007) Environmental and historical constraints on global patterns of amphibian richness. *Proc R Soc Lond B Biol Sci* 274:1167–1173. <https://doi.org/10.1098/rspb.2006.0436>
- Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a practical information-theoretic approach. Springer-Verlag, New York
- Calcagno V (2015) Package 'glmulti' version 1.0.7.. <https://cran.r-project.org/web/packages/glmulti>. Accessed 24 Sept 2017
- Calcagno V, de Mazancourt C (2010) Glmulti: an R package for easy automated model selection with (generalized) linear models. *J Stat Softw* 34:1–29
- Chander G, Markham BL, Helder DL (2009) Summary of current radiometric calibration coefficients for Landsat MSS, TM, ETM+, and EO-1 ALI sensors. *Remote Sens Environ* 113:893–903. <https://doi.org/10.1016/j.rse.2009.01.007>
- Cutler DR, Edwards TC Jr, Beard KH, Cutler A, Hess KT, Gibson J, Lawler JJ (2007) Random forests for classification in ecology. *Ecology* 88:2783–2792. <https://doi.org/10.1890/07-0539.1>
- Delaugerre M, Cheylan M (1992) Atlas de Répartition des Batraciens et Reptiles de Corse. Parc Naturel Régional de Corse-Ecole Pratique des Hautes Études, Ajaccio
- Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, Marquéz JRG, Gruber B, Lafourcade B, Leitão PJ, Münkemüller T (2013) Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36:27–46. <https://doi.org/10.1111/j.1600-0587.2012.07348.x>
- Duguet R, Melki F (2003) Les Amphibiens de France, Belgique et Luxembourg. Biotope, Mèze
- Escoriza D, Ben Hassine J (2017) Habitat structure and presence of *Salamandra algira* Bedriaga, 1883 in the Edough Peninsula, north-eastern Algeria. *Afr J Herpetol* 66:154–168. <https://doi.org/10.1080/21564574.2017.1388855>

- Escoriza D, Pascual G, Sánchez-Vialas A (2018) Habitat use in south-west European skinks (genus *Chalcides*). *PeerJ* 6:4274. <https://doi.org/10.7717/peerj.4274>
- Esselman PC, Allan JD (2011) Application of species distribution models and conservation planning software to the design of a reserve network for the riverine fishes of northeastern Mesoamerica. *Freshw Biol* 56:71–88. <https://doi.org/10.1111/j.1365-2427.2010.02417.x>
- Fick SE, Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int J Climatol* 37:4302–4315. <https://doi.org/10.1002/joc.5086>
- Freemark KE, Merriam HG (1986) Importance of area and habitat heterogeneity to bird assemblages in temperate forest fragments. *Biol Conserv* 36:115–141. [https://doi.org/10.1016/0006-3207\(86\)90002-9](https://doi.org/10.1016/0006-3207(86)90002-9)
- Gálhidy L, Mihók B, Hagyó A, Rajkai K, Standovár T (2006) Effects of gap size and associated changes in light and soil moisture on the understorey vegetation of a Hungarian beech forest. *Plant Ecol* 183:133–145. <https://doi.org/10.1007/s11258-005-9012-4>
- Gamisans J (2010) Le paysage végétal de la Corse. Albiana, Ajaccio
- Gamisans J (2014) Flore des maquis et des végétations associées de Corse. Albiana, Ajaccio
- Garden JG, Mcalpine CA, Possingham HP, Jones DN (2007) Habitat structure is more important than vegetation composition for local-level management of native terrestrial reptile and small mammal species living in urban remnants: a case study from Brisbane, Australia. *Austral Ecol* 32:669–685. <https://doi.org/10.1111/j.1442-9993.2007.01750.x>
- Giordano AR, Ridenhour BJ, Storfer A (2007) The influence of altitude and topography on genetic structure in the long-toed salamander (*Ambystoma macrodactylum*). *Mol Ecol* 16:1625–1637. <https://doi.org/10.1111/j.1365-294X.2006.03223.x>
- Godmann O, Kariş M, Göçmen B (2016) Geographic nestedness of *Lyciasalamandra billae* (Amphibia: Salamandridae) populations within *L. antalyana* and description of a new subspecies. *Zool Middle East* 62:255–260. <https://doi.org/10.1080/09397140.2016.1202902>
- Gómez-Rodríguez C, Díaz-Paniagua C, Bustamante J, Porthault A, Florencio M (2010) Inter-annual variability in amphibian assemblages: implications for diversity assessment and conservation. *Aquat Conserv: Mar Freshw Ecosyst* 20:668–677. <https://doi.org/10.1002/aqc.1134>
- Gregory RD, Willis SG, Jiguet F, Voříšek P, Klvaňová A, van Strien A, Huntley B, Collingham YC, Couvet D, Green RE (2009) An indicator of the impact of climatic change on European bird populations. *PLoS One* 4:4678. <https://doi.org/10.1371/journal.pone.0004678>
- Griffith MB, Husby P, Hall RK, Kaufmann PR, Hill BH (2003) Analysis of macroinvertebrate assemblages in relation to environmental gradients among lotic habitats of California's Central Valley. *Environ Monit Assess* 82: 281–309. <https://doi.org/10.1023/A:102199431>
- Guénette JS, Villard MA (2005) Thresholds in forest bird response to habitat alteration as quantitative targets for conservation. *Conserv Biol* 19:1168–1180. <https://doi.org/10.1111/j.1523-1739.2005.00085.x>
- Guo F, Wang G, Su Z, Liang H, Wang W, Lin F, Liu A (2016) What drives forest fire in Fujian, China? Evidence from logistic regression and random forests. *Intern J Wildland Fire* 25:505–519. <https://doi.org/10.1071/WF15121>
- Hastie T, Tibshirani R, Friedman J (2009) The elements of statistical learning. Springer-verlag, New York
- Hernandez A, Escoriza D, Hou M (2018) Patterns of niche diversification in south-east Asian crocodile newts. *Zool Anz* 276:86–93. <https://doi.org/10.1016/j.jcz.2018.06.001>
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *Int J Climatol* 25:1965–1978. <https://doi.org/10.1002/joc.1276>
- Huete A, Didan K, Miura T, Rodríguez EP, Gao X, Ferreira LG (2002) Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sens Environ* 83:195–213. [https://doi.org/10.1016/S0034-4257\(02\)00096-2](https://doi.org/10.1016/S0034-4257(02)00096-2)
- Hughes GO, Thuiller W, Midgley GF, Collins K (2008) Environmental change hastens the demise of the critically endangered riverine rabbit (*Bunolagus monticularis*). *Biol Conserv* 141:23–34. <https://doi.org/10.1016/j.biocon.2007.08.004>
- Hutchison VH (1961) Critical thermal maxima in salamanders. *Physiol Zool* 34:92–125
- IFN (Inventaire Forestier National) (2006) Inventaire forestier départemental Haute-Corse (2004) et Corse-du-Sud (2003): résultats du troisième inventaire. Ministère de l'Agriculture et de la Pêche, Montpellier
- IUCN (International Union for Conservation of Nature and Natural Resources) (2018) Red List of Threatened Species. <http://www.iucnredlist.org/>. Accessed 10 Aug 2018
- James FC, Wamer NO (1982) Relationships between temperate forest bird communities and vegetation structure. *Ecology* 63:159–171. <https://doi.org/10.2307/1937041>
- Jarvis A, Reuter HI, Nelson A, Guevara E (2008) Hole-filled SRTM for the globe Version 4. <http://www.cgiar-csi.org/data>. Accessed 6 May 2017
- Jeanmonod D, Gamisans J (2007) Flora Corsica. Édusud, Aix-en-Provence
- Johnston B, Frid L (2002) Clearcut logging restricts the movements of terrestrial Pacific giant salamanders (*Dicamptodon tenebrosus* Good). *Can J Zool* 80: 2170–2177. <https://doi.org/10.1139/z02-213>
- Kozak KH, Wiens JJ (2010) Niche conservatism drives elevational diversity patterns in Appalachian salamanders. *Am Nat* 176:40–54. <https://doi.org/10.1086/653031>
- Lalkhen AG, McCluskey A (2008) Clinical tests: sensitivity and specificity. *Contin Educ Anaesth Crit Care Pain* 8:221–223. <https://doi.org/10.1093/bjaceaccp/mkn041>
- Lanza B, Andreone F, Bologna MA, Corti C, Razzetti E (2007) Fauna d'Italia, Amphibia. Calderini, Bologna
- Larson DL, Anderson PJ, Newton W (2001) Alien plant invasion in mixed-grass prairie: effects of vegetation type and anthropogenic disturbance. *Ecol Appl* 11:128–141. [https://doi.org/10.1890/1051-0761\(2001\)011\[0128:APIIMGJ\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2001)011[0128:APIIMGJ]2.0.CO;2)
- Lescure J, de Massary JC (2012) Atlas des Amphibiens et Reptiles de France. Muséum National d'Histoire Naturelle, Paris
- Liaw A, Wiener M (2002) Classification and regression by randomForest. *R News* 2:18–22
- Loiselle BA, Howell CA, Graham CH, Goerck JM, Brooks T, Smith KG, Williams PH (2003) Avoiding pitfalls of using species distribution models in conservation planning. *Conserv Biol* 17:1591–1600. <https://doi.org/10.1111/j.1523-1739.2003.00233.x>
- Martel A, Spitzen-van der Sluijs A, Blooi M, Bert W, Ducatelle R, Fisher MC, Woeltjes A, Bosman W, Chiers K, Bossuyt F, Pasmans F (2013) *Batrachochytrium salamandrivorans* sp. nov. causes lethal chytridiomycosis in amphibians. *Proc Natl Acad Sci* 110:15325–15329. <https://doi.org/10.1073/pnas.1307356110>
- McIntyre PJ, Thorne JH, Dolanc CR, Flint AL, Flint LE, Kelly M, Ackerly DD (2015) Twentieth-century shifts in forest structure in California: denser forests, smaller trees, and increased dominance of oaks. *Proc Natl Acad Sci* 112: 1458–1463. <https://doi.org/10.1073/pnas.1410186112>
- Miaud C, Cheylan M, Sindaco R, Romano A (2009) *Salamandra corsica*. In: The IUCN red list of threatened species 2009: e.T59465A11926865. <https://doi.org/10.2305/IUCN.UK.2009.RLTS.T59465A11926865.en>. Accessed 15 Nov 2018
- Munwes J, Geffen E, Roll U, Friedmann A, Daya A, Tikochinski Y, Gafny S (2010) The change in genetic diversity down the core-edge gradient in the eastern spadefoot toad (*Pelobates syriacus*). *Mol Ecol* 19:2675–2689. <https://doi.org/10.1111/j.1365-294X.2010.04712.x>
- Mwitondi KS (2013) Data mining with rattle and R. *J Appl Stat* 40:464. <https://doi.org/10.1080/02664763.2012.749050>
- Peel MC, Finlayson BL, McMahon TA (2007) Updated world map of the Köppen-Geiger climate classification. *Hydrol Earth Syst Sci Discuss* 4:439–473
- Petranka JW, Eldridge ME, Haley KE (1993) Effects of timber harvesting on southern Appalachian salamanders. *Conserv Biol* 7:363–370. <https://doi.org/10.1046/j.1523-1739.1993.07020363.x>
- Pettorelli N, Vik JO, Mysterud A, Gaillard JM, Tucker CJ, Stenseth NC (2005) Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends Ecol Evol* 20:503–510. <https://doi.org/10.1016/j.tree.2005.05.011>
- QGIS Development Team (2018) Quantum GIS vs 3.4.1. Open Source Geospatial Foundation Project. <https://www.qgis.org>. Accessed 15 Nov 2018
- R Core Team (2018) R: A language and environment for statistical computing: vs 3.5.1. R Foundation for Statistical Computing, Vienna
- Rasband W (2018) ImageJ vs 1.52a. <https://imagej.nih.gov/ij>. Accessed 27 Sept 2018
- Riley SJ, DeGloria SD, Elliot R (1999) A terrain ruggedness index that quantifies topographic heterogeneity. *Interm J Sci* 5:23–27
- Roces-Díaz JV, Jiménez-Alfaro B, Chytrý M, Díaz-Varela ER, Álvarez-Álvarez P (2018) Glacial refugia and mid-Holocene expansion delineate the current distribution of *Castanea sativa* in Europe. *Palaeogeogr Palaeoclimatol Palaeoecol* 491:152–160. <https://doi.org/10.1016/j.palaeo.2017.12.004>
- Rödel MO, Ernst R (2004) Measuring and monitoring amphibian diversity in tropical forests. I. an evaluation of methods with recommendations for standardization. *Ecotropica* 10:1–14
- Rovito SM, Parra-Olea G, Vásquez-Almazán CR, Papenfuss TJ, Wake DB (2009) Dramatic declines in neotropical salamander populations are an important part of the global amphibian crisis. *Proc Natl Acad Sci* 106:3231–3236. <https://doi.org/10.1073/pnas.0813051106>

- Salvidio S (1998) Estimating abundance and biomass of a *Speleomantes strinatii* (Caudata, Plethodontidae) population by temporary removal sampling. *Amphibia-Reptilia* 19:113–124
- Sax DF, Gaines SD (2008) Species invasions and extinction: the future of native biodiversity on islands. *Proc Natl Acad Sci* 105:11490–11497. <https://doi.org/10.1073/pnas.0802290105>
- Scheffers BR, Edwards DP, Diesmos A, Williams SE, Evans TA (2014a) Microhabitats reduce animal's exposure to climate extremes. *Glob Change Biol* 20:495–503. <https://doi.org/10.1111/gcb.12439>
- Scheffers BR, Evans TA, Williams SE, Edwards DP (2014b) Microhabitats in the tropics buffer temperature in a globally coherent manner. *Biol Lett* 10: 20140819. <https://doi.org/10.1098/rsbl.2014.0819>
- Sing T, Sander O, Beerenwinkel N, Lengauer T (2015) Package 'ROCR' vs 1.0–7. <https://cran.r-project.org/web/packages/ROCR/index.html>. Accessed 2 Aug 2017
- Spotila JR (1972) Role of temperature and water in the ecology of lungless salamanders. *Ecol Monogr* 42:95–125. <https://doi.org/10.2307/1942232>
- Suggitt AJ, Gillingham PK, Hill JK, Huntley B, Kunin WE, Roy DB, Thomas CD (2011) Habitat microclimates drive fine-scale variation in extreme temperatures. *Oikos* 120:1–8. <https://doi.org/10.1111/j.1600-0706.2010.18270.x>
- Surasinghe TD, Baldwin RF (2015) Importance of riparian forest buffers in conservation of stream biodiversity: responses to land uses by stream-associated salamanders across two southeastern temperate ecoregions. *J Herpetol* 49:83–94
- Thorogood C (2018) *Guide des Fleurs Sauvages de Méditerranée Occidentale*. Delachaux et Niestlé, Paris
- van Geldern R, Kuhlemann J, Schiebel R, Taubald H, Barth JA (2014) Stable water isotope patterns in a climate change hotspot: the isotope hydrology framework of Corsica (western Mediterranean). *Isot Environ Health Stud* 50: 184–200. <https://doi.org/10.1080/10256016.2013.839559>
- von Arx G, Dobbertin M, Rebetez M (2012) Spatio-temporal effects of forest canopy on understory microclimate in a long-term experiment in Switzerland. *Agric For Meteorol* 166:144–155. <https://doi.org/10.1016/j.agrformet.2012.07.018>
- Warburg MR (1986) A relic population of *Salamandra salamandra* on Mt. Carmel: a ten-year study. *Herpetologica* 42:174–179
- Weiner J (2017) *pca3d*: three dimensional PCA Plots. R package version 0.10. <https://CRAN.R-project.org/package=pca3d>. Accessed 12 May 2016
- Williams AK, Berkson J (2004) Reducing false absences in survey data: detection probabilities of red-backed salamanders. *J Wildl Manag* 68:418–428. [https://doi.org/10.2193/0022-541X\(2004\)068\[0418:RFAISD\]2.0.CO;2](https://doi.org/10.2193/0022-541X(2004)068[0418:RFAISD]2.0.CO;2)
- Williams SE, Marsh H, Winter J (2002) Spatial scale, species diversity, and habitat structure: small mammals in Australian tropical rain forest. *Ecology* 83:1317–1329. [https://doi.org/10.1890/0012-9658\(2002\)083\[1317:SSSDAH\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[1317:SSSDAH]2.0.CO;2)
- Wilson SD, Keddy PA (1986) Species competitive ability and position along a natural stress/disturbance gradient. *Ecology* 67:1236–1242. <https://doi.org/10.2307/1938679>

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
